

Magnetic backgrounds and tachyonic instabilities in closed superstring theory and M-theory

J.G. Russo^{a,b} * and A.A. Tseytlin^c * †

^a *Departamento de Física, Universidad de Buenos Aires,
Ciudad Universitaria, Pab. I, 1428 Buenos Aires, and Conicet*

^b *The Abdus Salam International Center for Theoretical Physics,
Strada Costiera 11, 34014 Trieste, Italy.*

^c *Smith Laboratory, Ohio State University,
Columbus OH 43210-1106, USA*

Abstract

Models of closed superstrings in certain curved NS-NS magnetic flux backgrounds are exactly solvable in terms of free fields. They interpolate between free superstring theories with periodic and antiperiodic boundary conditions for fermions around some compact direction, and, in particular, between type 0 and type II string theories. Using “9–11” flip, this interpolation can be extended to M-theory and provides an interesting setting for study of tachyon problem in closed string theory. Starting with a general 2-parameter family of such Melvin-type models, we present several new magnetic flux backgrounds in 10-d string theory and 11-d M-theory and discuss their tachyonic instabilities. In particular, we suggest a description of type 0B theory in terms of M-theory in curved magnetic flux background, which supports its conjectured $SL(2, Z)$ symmetry, and in which the type 0 tachyon appears to correspond to a state in $d = 11$ supergravity multiplet. In another “T-dual” description, the tachyon is related to a winding membrane state.

April 2001

* e-mail address: russo@df.uba.ar

* e-mail address: tseytlin.1@osu.edu

† Also at Imperial College, London and Lebedev Physics Institute, Moscow.

1. Introduction

One of the fundamental problems in string theory is vacuum stability. Non-supersymmetric vacua are often unstable, containing tachyons. Recently, there was a progress in understanding the issue of open string tachyon condensation. The closed string tachyon problem is harder, and to try to address it, type 0 string theory [1] is a natural model to study. As was noted in [2], in contrast to what happens in the open string case, the type 0 string partition function in tachyon background does not seem to contain a tree-level potential for the tachyon, i.e. a stable vacuum cannot be determined just from analysis of low-energy tree-level effective theory in flat background.

There have been two proposals on tachyon stabilization in type 0 theory – one within string perturbation theory but in curved background, and another based on non-perturbative M-theory formulation:

- (i) it was suggested in [3,4] that type 0B string in $AdS_5 \times S^5$ + 5-form R-R background becomes stable for small enough scale of the space;¹
- (ii) it was conjectured in [7] that the tachyon of type 0A string in flat space gets $m^2 > 0$ at strong coupling (where type 0A string becomes dual to 11-d M-theory on large circle with antiperiodic fermions). Ref. [8] combined this second proposal with the observation that Kaluza-Klein Melvin [9] magnetic background [10,11,12,13] in type II string theory and in 11-d M-theory may be used to interpolate [14,15] between periodic and antiperiodic boundary conditions for space-time fermions. This suggests a dynamical mechanism of type 0 string tachyon stabilization – it disappears at strong magnetic R-R field [8,16].

Our aim will be to extend the discussion in [15,8] to the case of more general class of Melvin-type backgrounds with two independent magnetic parameters b and $\tilde{b}$. It includes K-K Melvin ($b \neq 0, \tilde{b} = 0$) and dilatonic Melvin ($b = \tilde{b}$) [10,17] solutions as special cases and is covariant under T-duality. The bosonic [18] and superstring [14,19] models in the

¹ This suggestion is motivated by the fact that R-R flux shifts the tachyon (mass)² in positive direction. To prove this conjecture one needs to demonstrate that α' -corrections do not spoil this effect, i.e. to solve the corresponding R-R sigma model. Using a generalization of AdS/CFT correspondence, for $\alpha'R \sim 1$ one may expect to get some input from the dual large N gauge theory description which is weakly coupled in that regime [5]. For a recent interesting discussion of issues related to closed string tachyons in orbifold theories in the context of AdS/CFT correspondence (and a suggestion that type 0 theory may still remain unstable at small radius due to a new tachyon appearing in the twisted sector) see [6].

corresponding NS-NS backgrounds are exactly solvable, with perturbative spectrum and torus partition function known explicitly (see [20] for a review).

Here we will lift these type IIA magnetic solutions to 11 dimensions, getting a $\tilde{b} \neq 0$ generalization of the flat [13,12] $d = 11$ background discussed in [15,8]. Dimensional reduction along different directions (or U-dualities directly in $d = 10$) lead to a number of $d = 10$ supergravity backgrounds with R-R magnetic fluxes, generalizing to $\tilde{b} \neq 0$ the R-R magnetic flux 7-brane [15,8] of type IIA theory. We shall investigate instabilities of these backgrounds.

Our motivation is to get a more detailed understanding of magnetic interpolations between stable and unstable theories, combining previous perturbative results [14] about spectra of NS-NS models with non-perturbative $d = 11$ extensions based, as in [7,8], on conjectured applicability of the “9-11” flip. One general aim is to establish precise relations between non-supersymmetric backgrounds in type II superstring theory and unstable backgrounds in type 0 theory. Another is to use interpolating magnetic backgrounds to connect D-brane solutions in the two theories, and, hopefully, shed more light on non-supersymmetric gauge theories dual to them. In the process, one may also learn more about various non-supersymmetric compactifications of M-theory.

Before describing the content of the paper let us recall some basic points about the relation between type 0 and type II theories. First, there is a perturbative relation in flat space: weakly coupled type 0 string theory may be interpreted as an orbifold of type II theory [21]. Indeed, type II theory compactified on a circle with *anti*periodic boundary conditions for space-time fermions is closely related to type 0 theory compactified on a circle [22,23]. More precisely, the two theories are limits of the interpolating “9-dimensional” theory [21] – Σ_R orbifold of type II theory. Σ_R stands for $(S^1)_R/((-1)^{F_s} \times \mathcal{S})$, where F_s is space-time fermion number and $\mathcal{S}$ is half-shift along the circle ($X_9 \rightarrow X_9 + \pi R$).² Type IIA on $\Sigma_{R \rightarrow \infty}$ is type IIA theory on an infinite circle and type IIA on $\Sigma_{R \rightarrow 0}$ is type 0A theory

² The role of $\mathcal{S}$ is to mix $(-1)^{F_s}$ orbifold with compactification on the circle, allowing for the interpolation [21]. Invariant states under $\mathcal{S}$ have even momentum quantum numbers m and integer winding numbers w , while twisted sector states have odd m and half-integer w . The action of $\mathcal{S}$ is irrelevant for $R \rightarrow \infty$, but is crucial for reproducing type 0 spectrum for $R \rightarrow 0$. The existence of this 9-d interpolating orbifold theory allows one to establish relations between D-branes in type II and type 0 theories [4,24].

on $(S^1)_{R \rightarrow 0}$ (or T-dual type 0B theory on $(S^1)_{R \rightarrow \infty}$). Thus type IIA on Σ interpolates between type IIA and type 0B *ten*-dimensional theories [21].³

Using that type II superstring in the $d = 10$ K-K Melvin NS-NS background $ds^2 = -dt^2 + dx_s^2 + dx_9^2 + dr^2 + r^2(d\varphi + b dx_9)^2$ (with $x_9 \equiv x_9 + 2\pi R$) is, for $bR = 1$, equivalent to type II theory on $R^9 \times (S^1)_R$ with antiperiodic boundary conditions for the space-time fermions along the x_9 circle [14],⁴ one may notice [8] that it is equivalent to the above orbifold of type II on $\Sigma_{R'}$ with the radius $R' = 2R$ (extra 2 is related to the shift $\mathcal{S}$).⁵ Therefore, this NS-NS Melvin model describes, in the limit $R \rightarrow 0$, weakly coupled type 0 string on $R^{1,8} \times (S^1)_{R \rightarrow 0}$ (or type 0B on $R^{1,8} \times (S^1)_{R \rightarrow \infty}$).⁶ To summarize, type II orbifold on Σ_{2R} is the same as type II theory in Melvin background at $bR = 1$ and thus the latter model interpolates between type II and type 0 theories in infinite $d = 10$ space.

This can be extended to a general claim [8] that type IIA(B) theory in the Kaluza-Klein NS-NS Melvin background with parameters (b, R) is equivalent (has the same perturbative spectrum and thus same torus partition function) to the orbifold of type IIA(B) on $\Sigma_{R'}$ in this NS-NS Melvin background with parameters $(b' = b - R^{-1}, R' = 2R)$.

³ In general, there are 4 different perturbative interpolating 9-d theories [21]: (a) type IIA on Σ_R (relating d=10 type IIA and type 0B); (b) type IIB on Σ_R (relating d=10 type IIB and type 0A); (c) type IIA on $(S^1)_R =$ type IIB on $(S^1)_{\tilde{R}}$, $\tilde{R} = \alpha'/R$ (relating d=10 type IIA and type IIB); (d) type 0A on $(S^1)_R =$ type 0B on $(S^1)_{\tilde{R}}$, $\tilde{R} = \alpha' R^{-1}$ (relating d=10 type 0A and type 0B). These theories are not equivalent – they share only respective boundary points. The Σ_R orbifold of type IIA(B) theory is, at the same time, equivalent (T-dual) to the $S^1/((-1)^{F_r} \times \mathcal{S})$ orbifold of type 0B(A) theory (F_r is right-moving world-sheet fermion number) [7]. These interpolating orbifold theories have, in contrast to type 0 theory, massive fermions in their spectrum.

⁴ This model at $bR = 1$ becomes essentially the same as a (non-compact 2-plane) limit of the twisted 3-torus model of [22] or a Wick rotation of the finite temperature superstring theory [23,25].

⁵ In the Melvin model with $bR = 1$ the shift $x_9 \rightarrow x_9 + 2\pi R$ is equivalent to 2π rotation in the 2-plane under which space-time fermions change sign. In the orbifold, the corresponding shift $\mathcal{S}$ is by $\pi R'$.

⁶ Combined with the results of [14], this observation allows one to describe type 0 theory in $R^{1,8} \times (S^1)_{R \rightarrow \infty}$ in light-cone Green-Schwarz formulation, where the action is quadratic in (standard, periodic in σ) fermions, but fermions are not decoupled from the radial coordinate of S^1 – their covariant derivative has flat but topologically non-trivial connection $\sim b dx_9$. Redefining the fermions to eliminate this connection makes them antiperiodic in σ in the odd winding sector [14,19].

Returning back to the case of flat background, the above compactification of type IIA theory on Σ_{R_9} corresponds to M-theory on $(S^1)_{R'_{11}} \times (S^1)_{R_9}/[(-1)^{F_s} \times \mathcal{S}]$, where R_9 and R'_{11} are taken to be small. Making the 9-11 flip⁷ one may then conjecture [7] that one gets an equivalent description of this theory as an ordinary S^1 (i.e. periodic fermions) compactification of the $d = 10$ theory obtained from M-theory on $\Sigma_{R'_{11}}$. Then type 0A theory may be interpreted as M-theory on $\Sigma_{R'_{11}}$, or, essentially, as M-theory on S^1 with radius $\frac{1}{2}R'_{11}$ with periodic/antiperiodic boundary conditions for bosons/fermions.⁸ Type 0B theory in $R^{1,9}$ is then M-theory on $T^2/[(-1)^{F_s} \times \mathcal{S}]$ in the limit of zero volume of the 2-torus [7], and it was argued in [7] that, like type IIB theory, this theory should be $SL(2, Z)$ symmetric.

Lifting the above NS-NS Melvin metric to 11 dimensions, replacing $x_9 \leftrightarrow x_{11}$ and reducing it down to 10 dimensions along x_{11} gives a non-supersymmetric R-R Melvin 7-brane in type IIA theory [13,15]. The dilaton $e^\phi = g_s(1 + b^2 r^2)^{3/4}$ grows away from the core, i.e. for large r the theory becomes effectively 11-dimensional [15].

Combining the above observations, it was suggested in [8] that type IIA and type 0A theories in such $d = 10$ R-R Melvin backgrounds⁹ are non-perturbatively dual to each other, with parameters related by $b_0 = b - R_{11}^{-1}$, $g_{s0} = \frac{R'_{11}}{2\sqrt{\alpha'}} = \frac{R_{11}}{\sqrt{\alpha'}}$. For example, for $bR_{11} = 1$ the $d = 11$ Melvin theory has antiperiodic fermions on the circle R_{11} , while, according to [7], type 0 theory is M-theory with antiperiodic fermions on circle $\frac{1}{2}R'_{11}$. That means, in particular, that starting with type 0 theory and increasing the value of the R-R magnetic parameter b_0 , the tachyon should disappear at strong enough magnetic field [8] when the theory becomes strongly coupled – its weakly-coupled description is in terms of stable weakly coupled type II theory in R-R Melvin background with small b .

This paper is organized as follows. In section 2 we review the 2-parameter $(b, \tilde{b})$ NS-NS Melvin type II superstring model of [14]. We describe the periodicity of the mass

⁷ This means exchanging the roles of the two circles, assuming that there is indeed an interpolating coordinate-invariant M-theory in 11 dimensions. Applicability of this flip remains of course a conjecture.

⁸ The factor of 1/2 is again due to the half-shift $\mathcal{S}$. It suggests that the same factor should be present in the relation between string coupling constants. However, it is not clear how to decide this unambiguously since the relation to perturbative type 0 theory based on 9-11 flip applies only in the limit of zero radius, i.e. at zero coupling.

⁹ Any bosonic solution of type II supergravity can be embedded into type 0 theory provided the fields of the twisted sector (tachyon and second set of R-R bosons) are set equal to zero [3].

spectrum in magnetic field parameters, and show how at special values of the parameters the spectrum reduce to that of the free superstring theory with fermions antiperiodic in spatial direction (reproducing, in particular, the perturbative type 0 spectrum in the $R \rightarrow 0$ limit). We also discuss tachyonic instabilities, and explain how they can be investigated using the effective field theory.

In section 3 we consider the U-dual 2-parameter R-R Melvin magnetic background, and find the corresponding 11-dimensional solution. This 11-d background is no longer locally flat, in contrast to the K-K Melvin case ($b \neq 0, \tilde{b} = 0$) considered in [15,8].

In section 4 we discuss two alternative representations of type 0A/0B string theories in terms of M-theory compactifications in the magnetic backgrounds, and study the appearance of instabilities as magnetic field is varied. The representation of type 0B theory in terms of M-theory in curved ($b = 0, \tilde{b} \neq 0$) magnetic background makes manifest its conjectured $SL(2, Z)$ symmetry. We also discuss the role of winding membrane states in reproducing type 0 tachyon from M-theory perspective, and comment on some implications for M-theory at finite temperature.

In section 5 we consider the equation for the tachyon of weakly-coupled type 0A string theory in the R-R Melvin background and demonstrate that, as in [3,4], the R-R flux shifts the value of tachyon (mass)² in positive direction. As we explain, this fact may be considered as an additional argument in favor of the the conjecture that the type 0A tachyon should disappear at sufficiently strong magnetic R-R field.

In section 6 we present a general class of $d = 11$ magnetic flux brane solutions with several independent field parameters, generalizing the background of section 3. Solutions representing magnetic fluxbranes superposed with ordinary D-brane backgrounds may dynamically relate stable and unstable string theory configurations with R-R charges, in particular, type IIA branes and type 0A branes. Such solutions that can be obtained from $d = 11$ flux branes superposed with charged M2-branes and M5-branes, are constructed in section 7.

In all of these cases, the magnetic fluxes are concentrated near $r = 0$ and vanish at $r = \infty$. In the Appendix, we describe a different solution with a *uniform* R-R magnetic field, representing rotating space-time dual to the NS-NS background in [26].

2. String models with NS-NS magnetic flux backgrounds

In this section we shall summarize the results of [14] for the type II superstring models which describe static magnetic flux tube backgrounds. The bosonic part of the corresponding conformal σ model has the Lagrangian

$$L = \partial_+ x_i \partial_- x_i + \partial_+ r \partial_- r + \frac{r^2}{1 + \tilde{b}^2 r^2} [\partial_+ \varphi + (b + \tilde{b}) \partial_+ x_9] [\partial_- \varphi + (b - \tilde{b}) \partial_- x_9] \\ + \partial_+ x_9 \partial_- x_9 + \mathcal{R} [\phi_0 - \frac{1}{2} \ln(1 + \tilde{b}^2 r^2)] . \quad (2.1)$$

Here $x_i = (t, x_s)$, $s = 1, \dots, 6$ are the free string coordinates, x_9 is a periodic coordinate of radius $R = R_9$ and φ is the angular coordinate with period 2π ($\mathcal{R}$ is proportional to a background world-sheet curvature). The constants b and $\tilde{b}$ are the magnetic field parameters. Explicitly, the curved 10-d background geometry has the following string-frame metric, NS-NS 2-form and the dilaton

$$ds^2 = -dt^2 + dx_s^2 + dx_9^2 + dr^2 + \frac{r^2}{1 + \tilde{b}^2 r^2} [d\varphi + (b + \tilde{b}) dx_9] [d\varphi + (b - \tilde{b}) dx_9] , \quad (2.2)$$

$$B_2 = \frac{\tilde{b} r^2}{1 + \tilde{b}^2 r^2} d\varphi \wedge dx_9 , \quad e^{2(\phi - \phi_0)} = \frac{1}{1 + \tilde{b}^2 r^2} , \quad (2.3)$$

i.e. may be interpreted as a 6-brane background in $d = 10$ type II string theory. The magnetic Melvin-type interpretation becomes apparent upon dimensional reduction in the x_9 -direction. The resulting solution of $d = 9$ supergravity

$$ds_9^2 = -dt^2 + dx_s^2 + dr^2 + r^2 f^{-1} \tilde{f}^{-1} d\varphi^2 , \quad (2.4)$$

$$\mathcal{A}_\varphi = b r^2 f^{-1} , \quad \mathcal{B}_\varphi = -\tilde{b} r^2 \tilde{f}^{-1} , \quad (2.5)$$

$$e^{2(\phi - \phi_0)} = \tilde{f}^{-1} , \quad e^{2\sigma} = f \tilde{f}^{-1} , \quad f \equiv 1 + b^2 r^2 , \quad \tilde{f} \equiv 1 + \tilde{b}^2 r^2 , \quad (2.6)$$

describes a magnetic flux tube universe with the magnetic vector $\mathcal{A}$ (coming from the metric) and the axial-vector $\mathcal{B}$ (coming from the 2-form) and the non-constant dilaton ϕ and K-K scalar σ . Note the $b \leftrightarrow \tilde{b}$ (T-duality) covariance of this background.

The special case of $\tilde{b} = 0$ corresponds to a locally flat 10-d metric (thus the model (2.1) with $\tilde{b} = 0$ can be formally obtained from the free string model by the shift $\varphi \rightarrow \varphi + b x_9$)

$$ds^2 = -dt^2 + dx_s^2 + dx_9^2 + dr^2 + r^2 (d\varphi + b dx_9)^2 , \quad (2.7)$$

but the 9-d geometry (2.4) is still curved. In the notation of [10], the model with $\tilde{b} = 0$ and arbitrary b corresponds to the “ $a = \sqrt{3}$ ” (Kaluza-Klein) Melvin model (i.e. Melvin solution in $d = 4$ Einstein gravity coupled to a Kaluza-Klein $U(1)$ gauge field), whereas the one with $\tilde{b} = b$ is the “ $a = 1$ ” (dilaton) Melvin model. The model with $b = 0, \tilde{b} \neq 0$ is T-dual to the $\tilde{b} = 0, b \neq 0$ model and, as follows from (2.2), has the following curved $d = 10$ metric

$$ds_{10}^2 = -dt^2 + dx_s^2 + dr^2 + \tilde{f}^{-1}(dx_9^2 + r^2 d\varphi^2) . \quad (2.8)$$

The $(R, \tilde{b}, b)$ model is T-dual ($x_9 \rightarrow \tilde{x}_9$) to $(\tilde{R}, b, \tilde{b})$ model, with

$$\tilde{R} = \frac{\alpha'}{R} . \quad (2.9)$$

Note that the $b = \tilde{b}$ ($a = 1$ dilaton Melvin) model with $R = \tilde{R}$ is the ‘self-dual’ point. In view of this T-duality relation, one may say that string models with $\tilde{b} > b$ are effectively equivalent to models with $\tilde{b} < b$. For fixed b inequivalent models are parametrized by $\tilde{b}$ in the interval $0 \leq \tilde{b} \leq b$ (with $a = \sqrt{3}$ and $a = 1$ Melvin thus being the boundary points).

For generic values of b and $\tilde{b}$ all of the type II supersymmetries are broken [11,14]. Supersymmetries are restored at the special values $bR = 2n_1$ and $\tilde{b}\tilde{R} = 2n_2$ ($n_i = 0, \pm 1, \dots$), where the conformal model describes the standard flat-space type II superstring theory (see below).

According to (2.7), a shift of x_9 by period of the circle $2\pi R$ implies rotation in the plane by angle $2\pi bR$. Thus in the special case of $bR = 2k + 1$ ($k = 0, \pm 1, \dots$) the metric becomes topologically trivial. However, the superstring theory on this background is still non-trivial (not equivalent to that of standard flat space) since space-time fermions change sign under 2π rotation in the plane. That means that the $bR = 1$ case (all models with $bR = 2k + 1$ are equivalent) describes superstrings with *antiperiodic* boundary condition in x_9 , just as in the twisted 3-torus model of [22] or in the finite temperature case [23] (see also [27]). As was already discussed in section 1, the type IIA string in the space (2.7) with $bR = 1$ is equivalent, for $R \rightarrow 0$, to type 0A string in $R^{1,8} \times S_{R \rightarrow 0}^1$ or type 0B string in $R^{1,8} \times S_{R \rightarrow \infty}^1$. The tachyon of type 0 theory originates from a particular winding mode present in the $bR = 1$ type II model spectrum (see section 2.1).

2.1. String spectrum

Despite the fact that the σ -model (2.1) (supplemented with fermionic terms given in [14]) looks non-linear and represents a curved 10-d background (2.2), the corresponding superstring theory is exactly solvable in terms of free fields [14]. Using operator quantization the resulting string Hamiltonian and the level matching condition are given by [14]

$$\begin{aligned} \hat{H} = & \frac{1}{2}\alpha'(-E^2 + p_s^2) + \hat{N}_R + \hat{N}_L + \frac{1}{2}\tilde{R}R^{-1}(m - bR\hat{J})^2 \\ & + \frac{1}{2}R\tilde{R}^{-1}(w - \tilde{b}\tilde{R}\hat{J})^2 - \gamma(\hat{J}_R - \hat{J}_L) = 0 , \end{aligned} \quad (2.10)$$

$$\hat{N}_R - \hat{N}_L = mw , \quad (2.11)$$

where

$$\gamma \equiv bRw + \tilde{b}\tilde{R}m - bR\tilde{b}\tilde{R}\hat{J} . \quad (2.12)$$

Here p_s are continuous momenta in the 6 free directions, and the integers m and w represent quantized momentum and winding numbers in the compact x_9 direction.

$\hat{N}_R$ and $\hat{N}_L$ are the number of states operators, which have the standard free string theory form in terms of normal-ordered bilinears of bosonic and fermionic oscillators,

$$\hat{N}_{R,L} = N_{R,L} - a , \quad a^{(R)} = 0 , \quad a^{(\text{NS})} = \frac{1}{2} . \quad (2.13)$$

$\hat{J} \equiv \hat{J}_L + \hat{J}_R$ and $\hat{J}_{L,R}$ are the angular momentum operators in the 2-plane (i.e. constructed out of oscillators corresponding to the 2-plane coordinates $X = x_1 + ix_2 = re^{i\varphi}$) with the eigenvalues

$$\hat{J}_{L,R} = \pm(l_{L,R} + \frac{1}{2}) + S_{L,R} , \quad \hat{J} = l_L - l_R + S_L + S_R . \quad (2.14)$$

The orbital momenta $l_{L,R} = 0, 1, 2, \dots$ (which replace the continuous linear momenta p_1, p_2 in the (r, φ) 2-plane for non-zero values of γ) are the analogues of the Landau quantum number, and $S_{R,L}$ are the spin components (in the NS-NS sector, $|S_{R,L}| \leq \hat{N}_{R,L} + 1$). In the special case of $\gamma = 0$ (or, more generally, $\gamma = n$) the zero-mode structure changes in that the translational invariance in the 2-plane is restored. This leads to a modification in the above formulas, with the orbital momentum parts becoming the standard flat-space theory expressions in terms of the zero-mode operators $x_{1,2}, p_{1,2}$.

The expression for the Hamiltonian (2.10) applies for $-1 < \gamma < 1$, while for other values one is to replace γ by $\hat{\gamma} \equiv \gamma - [\gamma]$, where $[\dots]$ denotes the integer part.¹⁰ The mass spectrum is then invariant under

$$b \rightarrow b + 2n_1 R^{-1} , \quad \tilde{b} \rightarrow \tilde{b} + 2n_2 \tilde{R}^{-1} , \quad n_1, n_2 = 0, \pm 1, \dots \quad (2.15)$$

This becomes clear by observing that this transformation may be compensated by integer shifts of quantum numbers. Note that because $\hat{J}$ can take both integer (in NS-NS, R-R sectors) and *half*-integer (in NS-R, R-NS sectors) values, the symmetry of the bosonic part of the spectrum $b \rightarrow b + k_1 R^{-1}$, $\tilde{b} \rightarrow \tilde{b} + k_2 \tilde{R}^{-1}$ is *not* a symmetry of its fermionic part, i.e. the full superstring spectrum is invariant under (2.15) with even integer shifts $k_i = 2n_i$ only.

In the case of $bR = 2n_1$, $\tilde{b}\tilde{R} = 2n_2$ (i.e. $\gamma = \text{even integer}$, $\hat{\gamma} = 0$) the spectrum is thus equivalent to that of the standard free superstring compactified on a circle. In the two cases

$$(a) \ bR = \text{odd}, \ \tilde{b}\tilde{R} = \text{even} \quad \text{or} \quad (b) \ bR = \text{even}, \ \tilde{b}\tilde{R} = \text{odd} , \quad (2.16)$$

the spectrum is the same as that of the free superstring compactified on a circle with antiperiodic boundary conditions for space-time fermions [22,23].¹¹

The T-duality symmetry in the compact Kaluza-Klein direction x_9 (which exchanges the axial-vector and vector magnetic field parameters $\tilde{b}$ and b , cf. (2.5)) is manifest in (2.10): the Hamiltonian is invariant under $R \leftrightarrow \tilde{R} \equiv \alpha' R^{-1}$, $b \leftrightarrow \tilde{b}$, $m \leftrightarrow w$.

A close examination of (2.10) shows that all states with $\hat{J}_R - \hat{J}_L \leq \hat{N}_R + \hat{N}_L$ have positive (mass)². The only bosonic states which may be *tachyonic* thus lie on the first Regge trajectory with maximal value of S_R , minimal value of S_L , and zero orbital momentum,

¹⁰ This follows from a change in the normal ordering constant, see [18].

¹¹ This relation becomes particularly clear in the light-cone GS formulation given in [14]. For example, for $\tilde{b} = 0$ the GS action is quadratic in fermionic 2-d fields which are coupled to a flat but topologically non-trivial connection. After a formal redefinition, one gets free action in terms of fermion fields obeying the boundary condition $\Sigma_{R,L}(\tau, \sigma + \pi) = e^{i\pi\gamma} \Sigma_{R,L}(\tau, \sigma)$. This makes it clear that the inequivalent values of γ are $-1 < \gamma \leq 1$, with $\gamma = 1$ corresponding to the antiperiodic case. More precisely, the fermions are antiperiodic with respect to x_9 , but as functions of the string world-sheet direction σ their periodicity is determined by $\gamma = bRw$, i.e. they are periodic in σ in the even winding sector and antiperiodic in σ in the odd winding sector.

i.e. $\hat{J}_R = S_R - \frac{1}{2} = \hat{N}_R + \frac{1}{2}$, $\hat{J}_L = S_L + \frac{1}{2} = -\hat{N}_L - \frac{1}{2}$, so that $\hat{J}_R - \hat{J}_L = \hat{N}_R + \hat{N}_L + 1$. In particular, all fermionic states have $(\text{mass})^2 \geq 0$, as expected in a unitary theory.

From eq. (2.10) one learns that in general there are instabilities (associated with states with high spin and charge) for arbitrarily small values of the magnetic field parameters. The only exception is the $\tilde{b} = 0$ model (or its T-dual $b = 0$ model): here the type II superstring model has no tachyons if the value of b is smaller than some *finite* critical value b_{cr} . The spectrum of the $\tilde{b} = 0$ model is given by ($M^2 = E^2 - p_s^2$)

$$\alpha' M^2 = 2(\hat{N}_R + \hat{N}_L) + \alpha'(mR^{-1} - b\hat{J})^2 + \frac{w^2 R^2}{\alpha'} - 2\hat{\gamma}(\hat{J}_R - \hat{J}_L) , \quad (2.17)$$

$$\hat{\gamma} = \gamma - [\gamma] , \quad \gamma = bRw . \quad (2.18)$$

Here the first (lowest mass) potentially tachyonic state that appears as b is increased from zero has

$$\hat{N}_R = \hat{N}_L = 0 , \quad \hat{J} = 0 , \quad \hat{J}_R - \hat{J}_L = 1 , \quad m = 0 , \quad w = 1 , \quad (2.19)$$

$$l_R = l_L = 0 , \quad S_R = 1 , \quad S_L = -1 ,$$

so that (for $bR < 1$)

$$\alpha' M^2 = \frac{R^2}{\alpha'} - 2bR \equiv 2R(b_{\text{cr}} - b) , \quad b_{\text{cr}} = \frac{R}{2\alpha'} . \quad (2.20)$$

It becomes tachyonic for $b > b_{\text{cr}}$. Because of the periodicity of the spectrum (2.15) (cf. (2.12),(2.10) for $\tilde{b} = 0$) the parameter bR may be restricted to the interval

$$-1 < bR \leq 1 . \quad (2.21)$$

Then the stability condition $b < b_{\text{cr}}$ implies that the theory with

$$R \geq R_{\text{cr}} \equiv \sqrt{2\alpha'} \quad (2.22)$$

is tachyon free.

In (2.20) we assumed that $bR < 1$. For $bR = 1$, one gets $\hat{\gamma} = 0$ so that the last gyromagnetic term in (2.17) and thus the $-2bR$ term in (2.20) is absent. However, the spectrum is continuous at this point: there is an extra negative term coming from a change in values of normal ordering constants in $\hat{N}_R$ and $\hat{N}_L$ [14], which produce extra -2 term in $\alpha' M^2$ in (2.20) (see also below).

Similar considerations apply to the T-dual model with $b = 0$, $\tilde{b} \neq 0$, which has the equivalent spectrum. For this model there are tachyons for any $\tilde{b} > \tilde{b}_{\text{cr}}$, with $\tilde{b}_{\text{cr}} = \frac{\tilde{R}}{2\alpha'} = \frac{1}{2\tilde{R}}$. If we consider the region $0 \leq \tilde{b}\tilde{R} < 1$ then this model is tachyon-free if $\tilde{R} \geq \sqrt{2\alpha'}$, i.e. $R \leq \sqrt{\frac{\alpha'}{2}}$.

In order to interpret the spectrum which follows from (2.10) in terms of gyromagnetic interaction, i.e. to make explicit how the momenta $\hat{J}_R$ and $\hat{J}_L$ couple to the magnetic field, it is convenient to introduce the “left” and “right” magnetic field parameters $B_{L,R} \equiv b \pm \tilde{b}$ and charges $Q_{L,R} = mR^{-1} \pm w\tilde{R}^{-1}$. The mass formula (2.10) then takes the form

$$M^2 = M_{\text{free}}^2 - 2(B_L Q_L \hat{J}_R + B_R Q_R \hat{J}_L) + (B_L^2 \hat{J}_R + B_R^2 \hat{J}_L) \hat{J} , \quad (2.23)$$

$$M_{\text{free}}^2 \equiv \frac{2}{\alpha'} (\hat{N}_R + \hat{N}_L) + \frac{1}{2} (Q_L^2 + Q_R^2) . \quad (2.24)$$

In the model with $\tilde{b} = 0$ one has only the vector field background (see (2.5)), i.e. $B_L = B_R$. Then, as is clear from (2.23), the gyromagnetic coupling of the winding states involves the combination $\hat{J}_R - \hat{J}_L$, while the coupling of the momentum states – the combination $\hat{J}_R + \hat{J}_L$. In the T-dual model with $b = 0$ one has the axial-vector field background, $B_L = -B_R$, so that the momentum states couple to $\hat{J}_R - \hat{J}_L$ and the winding states to $\hat{J}_R + \hat{J}_L$. In both cases, the states which first become tachyonic have gyromagnetic coupling involving $\hat{J}_R - \hat{J}_L$.

The partition function of the model found in [14] exhibit the same symmetries as the spectrum:

$$Z(R, b, \tilde{b}) = Z(\tilde{R}, \tilde{b}, b) , \quad (2.25)$$

$$Z(R, b, \tilde{b}) = Z(R, b + 2n_1 R^{-1}, \tilde{b} + 2n_2 \tilde{R}^{-1}) , \quad n_{1,2} = 0, \pm 1, \dots , \quad (2.26)$$

where $\tilde{R}$ is the T-dual radius (2.9). Eq. (2.25) expresses the T-duality, while (2.26) is the symmetry (2.15) of the full conformal field theory.

In the special cases when both bR and $\tilde{b}\tilde{R}$ are even, the partition function is identically zero, since for these values the theory is equivalent to the free superstring theory with $b = \tilde{b} = 0$. Two other simple models which are equivalent to the free superstring theory with fermions obeying *antiperiodic* boundary conditions in the x_9 direction have parameters given in (2.16). For non-integer bR and $\tilde{b}\tilde{R}$ there are tachyons for any value of the radius R , and so the partition function contains infrared divergences. When $\tilde{b}\tilde{R}$ is equal to an even integer, Z in (2.26) reduces to that of the $\tilde{b} = 0$ (K-K Melvin) model, and is finite in the range of parameters for which there are no tachyons in the spectrum. Similar statement

applies in the T-dual case of $bR = 2n$. When either $\tilde{b}\tilde{R} = \alpha'\tilde{b}R^{-1}$ or bR is an odd number, the partition function is also finite in a certain range of values of the radius.

As was already mentioned above, the type IIA(B) theory in the background (2.7) or (2.2),(2.3) with $bR = 1, \tilde{b} = 0$ is equivalent to the orbifold of type IIA(B) on $(S^1)_{2R}/[(-1)^{F_s} \times \mathcal{S}]$, i.e. to the theory interpolating between type IIA(B) in $R^{1,8} \times (S^1)_{R \rightarrow \infty}$ and type 0A(B) in $R^{1,8} \times (S^1)_{R \rightarrow 0}$. In this case $\hat{\gamma} = 0$, and the spectrum (2.17) coincides with the spectrum of free superstring theory with antiperiodic fermions. As well known, in this case tachyons may appear in the $w = \text{odd}$ sector, because of the reversal of the GSO condition [23]. The scalar state with winding number $w = 1$ and $N_R = N_L = 0$ (i.e. with $\hat{N}_R = \hat{N}_L = -\frac{1}{2}$)¹² is not projected out by the GSO, and has the mass¹³

$$M^2 = \frac{R^2}{\alpha'^2} - \frac{2}{\alpha'} . \quad (2.27)$$

In the limit of $R \rightarrow 0$, it represents the only tachyonic state that is present in the spectrum of type 0B(A) theory in $R^{1,9}$. It is interesting to see how this negative mass shift emerges as the magnetic field is increased from zero. The tachyon corresponds to the state (2.19), whose mass (2.20), for $\gamma = bR = 1 - \epsilon = \hat{\gamma}$, reduces to (2.27) in the limit $\epsilon \rightarrow 0$. In that sense type 0 tachyon corresponds to the special “spinning” winding mode of type II theory in the background (2.7), and its mass $-\frac{2}{\alpha'}$ has its origin in the gyromagnetic interaction term.

Let us note in passing that some of the expressions for the mode expansions and the spectrum in this magnetic model (see [26,18,14]) formally are very similar to the corresponding ones for the open strings in the presence of a constant magnetic (F_{mn} or B_{mn}) field [28]. One may think, by analogy with the open string case [29], that one can then use certain limits of these magnetic flux tube models to relate *closed* strings to non-commutative field theories. However, this does not seem to be possible. The point is that open strings in constant NS-NS B_{mn} field behave as electric dipoles with total charge equal to zero, so that there is a residual translational invariance and non-zero commutators for the center of mass coordinates. In contrast, closed string states can have only charges associated with momentum or winding number, i.e. they do not behave like electric dipoles.

¹² This is transparent in the GS formulation: in the odd winding (antiperiodic) sector the operators $N_{L,R}$ have half-integer eigenvalues and the normal-ordering constants are equal to $-1/2$.

¹³ The extra (twisted sector) set of R-R massless states of type 0 theory also originate from the odd winding sector with $\hat{N}_R = \hat{N}_L = 0$.

For charged closed strings with $\gamma \neq n$, there is no translational invariance in the x_1, x_2 plane (there are Landau levels instead) and there is no obvious analog of the non-zero commutator $[x_1, x_2]$ which appears in the open string case (one may introduce operators x_1, x_2 formally defined in terms of Landau level creation/annihilation operators, which in the zero field limit reduce to the standard center of mass coordinates, but they commute).

2.2. Tachyon spectrum from effective field theory

The full string mass spectrum (2.10) contains, in particular, the spectrum of small fluctuations of type II supergravity fields expanded near the curved background (2.2),(2.3). To decouple string modes one is to take $\alpha' \rightarrow 0$ (assuming that $\sqrt{\alpha'}/R, \sqrt{\alpha'}b, \sqrt{\alpha'}\tilde{b} \rightarrow 0$).¹⁴

In the model with $b = 0$ (2.8) the first state that becomes tachyonic as the field $\tilde{b}$ is increased from zero is from the “massless” sector of the theory, i.e. it has $\hat{N}_R = \hat{N}_L = 0$ and $w = 0$, $S_R = 1$, $S_L = -1$ (cf. eq. (2.19) in T-dual theory). This implies that one should be able to detect this instability directly from the effective supergravity field equations, i.e. without solving first the conformal string model. It is useful to see how this happens explicitly, since similar supergravity-based considerations may apply in other related cases (magnetic R-R backgrounds, or magnetic backgrounds in $d = 11$) where one does not have the benefit of knowing the exact microscopic (string theory or M-theory) spectrum.

Consider the general magnetic flux tube model (2.2),(2.3) with both $b, \tilde{b} \neq 0$. In the sector $\hat{N}_R = \hat{N}_L = 0$, $w = 0$ the mass formula (2.10) becomes

$$M^2 = \left(\frac{m}{R} - b\hat{J}\right)^2 + \tilde{b}^2\hat{J}^2 + 2\tilde{b}\left(\frac{m}{R} - b\hat{J}\right)(l_R + l_L + 1 - S_R + S_L) , \quad (2.28)$$

$$\hat{J} = l_L - l_R + S_L + S_R , \quad -1 \leq S_{R,L} \leq 1 .$$

Assuming $\tilde{b}(\frac{m}{R} - b\hat{J}) = \gamma/\alpha' > 0$ (see (2.12)), M^2 can become negative, provided $S_R = 1$, $S_L = -1$, and $l_L = l_R = 0$.

¹⁴ Since there is a compact x_9 direction, one may try to go further and consider, in addition to $\alpha' \rightarrow 0$, the low-energy approximation that ignores not only the massive string modes but also the Kaluza-Klein modes. Let us note that the Kaluza-Klein truncation is *not* a valid approximation near $bR = k_1$, or $\tilde{b}\tilde{R} = k_2$ (for integer k_1, k_2), where the theory is essentially ten-dimensional. This can be seen from the above mass spectrum (2.10),(2.23). For example, consider the case of $\tilde{b} = 0$ and the non-winding supergravity sector ($\hat{N}_R = \hat{N}_L = 0$, $w = 0$), so that $\gamma = 0$ (see (2.12)). Then the mass formula (2.17) becomes $M = R^{-1}(m - bR\hat{J})$, so that when bR is near 1, there are Kaluza-Klein states which become light. These are the states with $m = \hat{J}$ which have mass $M = \frac{m}{R_{\text{eff}}}$, $R_{\text{eff}} \equiv \frac{R}{1-bR}$. The Kaluza-Klein approximation thus fails when R_{eff} is large.

Let us now try to reproduce this spectrum of instabilities from the effective field theory. It is useful to start with a scalar supergravity fluctuation mode ψ that satisfies the “massless” wave equation in the background (2.2),(2.3),

$$\partial_m(e^{-2\phi}\sqrt{G}G^{mn}\partial_n)\psi = 0 . \quad (2.29)$$

Such spinless mode is not expected to become tachyonic since, according to (2.28), the tachyonic state should have a non-zero spin, but we shall see that (2.29) does reproduce the expected spectrum (2.28) in the sector $S_R = S_L = 0$.

Using (2.2),(2.3), eq. (2.29) takes the form

$$\begin{aligned} & \left[-\partial_t^2 + \partial_{x_s}^2 + \frac{1}{r}\partial_r(r\partial_r) + \frac{1}{r^2}(1+b^2r^2)(1+\tilde{b}^2r^2)\partial_\varphi^2 \right. \\ & \left. + (1+\tilde{b}^2r^2)\partial_{x_9}^2 - 2b(1+\tilde{b}^2r^2)\partial_\varphi\partial_{x_9} \right] \psi = 0 . \end{aligned} \quad (2.30)$$

Parametrizing the eigen-mode as

$$\psi = e^{iEt+ip_sx_s+ip_9x_9+il\varphi}\eta(r) , \quad p_9 = \frac{m}{R} , \quad m, l = 0, \pm 1, \dots , \quad (2.31)$$

we get from (2.30)

$$\left[-\frac{1}{r}\partial_r(r\partial_r) + \frac{l^2}{r^2} + \nu^2r^2 \right] \eta(r) = \mu^2\eta(r) , \quad (2.32)$$

where

$$\nu = \tilde{b}(p_9 - bl) , \quad \mu^2 = M^2 - (p_9 - bl)^2 - \tilde{b}^2l^2 , \quad M^2 \equiv E^2 - p_s^2 . \quad (2.33)$$

This is the Schrödinger equation for a two-dimensional oscillator with the frequency ν . It has normalizable solutions only if $\mu^2 = 2\nu(l_L + l_R + 1)$, i.e.

$$M_{\text{spin } 0}^2 = (p_9 - bl)^2 + \tilde{b}^2l^2 + 2\tilde{b}(p_9 - bl)(l_L + l_R + 1) , \quad l_L, l_R = 0, 1, 2, \dots . \quad (2.34)$$

Here l_L, l_R are related to the Landau level l and the radial quantum number k by

$$l_L - l_R = l , \quad l_L + l_R = 2k + |l| . \quad (2.35)$$

The mass spectrum (2.34) coincides indeed with (2.28) after one sets there $S_R = S_L = 0$, as appropriate for a scalar fluctuation. This scalar mode has $M^2 \geq 0$.

The mode that becomes tachyonic should have, according to (2.28), the non-vanishing spin in the 2-plane, $S_R = 1, S_L = -1$. Deriving the analog of eqs. (2.29),(2.30) for such

fluctuation directly from the supergravity equations is somewhat complicated.¹⁵ We shall by-pass this step by making a natural conjecture that to generalize (2.34) to the case of non-zero spin of the fluctuation one is simply to add the spins $S_{L,R}$ to the orbital momenta $l_{L,R}$ in (2.34). This conjecture is indeed confirmed by the expression (2.28) following from string theory.

In the case of interest ($S_R = -S_L = 1$) we obtain

$$M^2 = (p_9 - bl)^2 + \tilde{b}^2 l^2 + 2\tilde{b}(p_9 - bl)(l_L + l_R - 1) , \quad l_L, l_R = 0, 1, 2, \dots \quad (2.36)$$

As was already mentioned above, and as is now clear from (2.36), the first tachyon that appears for $\tilde{b} > \tilde{b}_{\text{cr}}$ ($\tilde{b}_{\text{cr}} = \frac{1}{2R}$) is the state with $l_R = l_L = 0$.

Let us note that in addition to the perturbative instability discussed above, the magnetic backgrounds in *supergravity* suffer also from a non-perturbative instability – a decay into monopole-antimonopole pairs [11,12]. This instability is present for any value of the magnetic field, i.e. even when perturbative instability is absent.

3. $d = 10$ supergravity backgrounds with Ramond-Ramond magnetic fluxes and their $d = 11$ counterparts

By using U-dualities, it is easy to convert the NS-NS magnetic background (2.2),(2.3) into $d = 10$ type II supergravity backgrounds with magnetic fluxes from the R-R sector. For example, starting with (2.2),(2.3) with $\tilde{b} \neq 0$ (i.e. with non-vanishing 2-form field) considered as solution of type IIB supergravity and applying S-duality we obtain another solution with non-vanishing R-R 2-form field (i.e. with the R-R axial-vector field in $d = 9$, cf. (2.5)).

In contrast to the NS-NS model (2.1), it is not clear how to solve the corresponding R-R superstring models: the Green-Schwarz action in the light-cone gauge [15] remains non-linear and is not related to a free theory by T-dualities and coordinate shifts, as was the case for (2.1) [14].

¹⁵ The unstable mode should be a particular mixture of the metric and the antisymmetric 2-form perturbations. There is some analogy with non-abelian gauge theory where the tachyon mode in a constant magnetic background has spin 1 [30].

3.1. Magnetic flux 7-brane of type IIA theory

Another example of R-R magnetic background is provided by the type IIA solution related to the $\tilde{b} = 0$ NS-NS Melvin background (2.2),(2.3) of type IIA theory by the chain of duality transformations: $T_{x_9}ST_{x_9}$. This gives the following background which may be interpreted as a non-supersymmetric R-R magnetic flux 7-brane of type IIA theory [15]

$$ds_{10}^2 = f^{1/2}(-dt^2 + dx_s^2 + dx_9^2 + dr^2 + r^2 f^{-1} d\varphi^2) , \quad (3.1)$$

$$e^{2\phi} = e^{2\phi_0} f^{3/2} , \quad A_\varphi = br^2 f^{-1} , \quad f \equiv 1 + b^2 r^2 , \quad (3.2)$$

where A is the R-R 1-form with the field strength

$$F = \frac{2b}{(1 + b^2 r^2)^2} r dr \wedge d\varphi = 2bf^{-2} e^r \wedge e^\varphi , \quad (3.3)$$

which is approximately constant near the core. The 7-brane directions are $x_s (s = 1, \dots, 6)$ and x_9 . Like the $d = 9$ reduction of K-K Melvin background [12], this solution can be obtained also directly [15] by dimensional reduction of the $d = 11$ background with trivial 3-form field $C_{\mu\nu\rho} = 0$ and the metric (cf. (2.7)) [12]

$$ds_{11}^2 = -dt^2 + dx_s^2 + dx_9^2 + dr^2 + r^2 (d\varphi + b dx_{11})^2 + dx_{11}^2 . \quad (3.4)$$

Here $\varphi \equiv \varphi + 2\pi$ and x_{11} has period $2\pi R_{11}$, so this metric is topologically non-trivial if $bR_{11} \neq n$. Since the metric is locally flat this is an exact solution to the equations of motion of $d = 11$ supergravity *and* M-theory (all possible higher-order curvature corrections vanish).

If the $11 \rightarrow 10$ dimensional reduction is done along the coordinate x_9 one obtains the $d = 10$ NS-NS background (2.2),(2.3) with $\tilde{b} = 0$ or (2.7) (K-K Melvin), discussed in [12,13]. Dimensional reduction along x_{11} gives the above type IIA solution (3.1),(3.2) with the magnetic R-R vector field. The direct $d = 11$ lift of the NS-NS background (2.7) is related to (3.4) by the 9-11 flip:

$$ds_{11}^2 = -dt^2 + dx_s^2 + dx_9^2 + dr^2 + r^2 (d\varphi + b dx_9)^2 + dx_{11}^2 . \quad (3.5)$$

If the effective string coupling e^ϕ in (3.2) is chosen to be small at $br \ll 1$ ($e^{\phi_0} \ll 1$) it gets large at $br \gg 1$, so one may say that the geometry is ten-dimensional close to the core of the 7-brane (inside the flux tube) but becomes effectively eleven-dimensional far from it [15,8].

The remarkable circumstance that the R-R 7-brane (3.1),(3.2) is a dimensional reduction of locally-flat $d = 11$ background (3.4) was utilized in [15] to derive the exact expression for the corresponding $d = 10$ GS superstring action starting with the known flat-space $d = 11$ supermembrane action [31].

It is interesting to note a similarity between the flux 7-brane (3.1),(3.2) and D6-brane of type IIA theory – both backgrounds have only the dilaton and the magnetic R-R vector as non-trivial matter fields.¹⁶ The dependence on the function $f(r)$ in the metric (3.1) and the dilaton field (3.2) is the same as the dependence on the inverse of the harmonic function $H^{-1} = (1 + \frac{L}{\rho})^{-1}$ in the D6-brane solution. This formal correspondence implies that applying T-duality along the “parallel” directions $(x_1, \dots, x_6, x_9)$ of the 7-brane one obtains similar flux p-branes having the same type of R-R fields and the same structure of f -dependence as in the Dp-brane solutions of IIA/B theories. In particular, we get the following type IIB flux 4-brane solution, which, like the D3-brane background, has *constant* dilaton. It has the metric

$$ds_{10}^2 = f^{1/2}(-dt^2 + dx_1^2 + \dots + dx_4^2 + dr^2) + f^{-1/2}(dx_5^2 + dx_6^2 + dx_9^2 + r^2 d\varphi^2) , \quad (3.6)$$

and the self-dual R-R 5-form with magnetic component $F_{r\varphi 569} \sim b f^{-2}$.

3.2. “Mixed” type IIA 2-parameter magnetic background

Starting with the general NS-NS solution (2.2),(2.3) and applying again $T_{x_9}ST_{x_9}$ dualities one obtains a generalization of the R-R Melvin solution (3.6), (3.2) in type IIA theory. It depends on two parameters which control the strengths of the R-R magnetic 1-form field A and the NS-NS 2-form field B_2 :

$$ds_{10A}^2 = f^{1/2}(-dt^2 + dx_s^2 + \tilde{f}^{-1}dx_9^2 + dr^2 + f^{-1}\tilde{f}^{-1}r^2 d\varphi^2) , \quad (3.7)$$

$$A = br^2 f^{-1} d\varphi , \quad B_2 = \tilde{b}r^2 \tilde{f}^{-1} d\varphi \wedge dx_9 , \quad e^{2\phi} = e^{2\phi_0} f^{3/2} \tilde{f}^{-1} , \quad (3.8)$$

$$f = 1 + b^2 r^2 , \quad \tilde{f} = 1 + \tilde{b}^2 r^2 .$$

The R-R 7-brane background (3.6),(3.2) is the special case of $\tilde{b} = 0$.

¹⁶ In a sense, this flux-brane may be viewed as “connecting” two magnetic R-R charges – D6-brane and anti D6-brane (cf. [13,8]).

Lifting this type IIA background to $d = 11$ we find the following “6-brane” solution of $d = 11$ supergravity

$$ds_{11}^2 = \tilde{f}^{1/3}(-dt^2 + dx_s^2) + \tilde{f}^{-2/3}[dx_9^2 + \tilde{f}dr^2 + r^2\tilde{f}^{-1}d\varphi^2 + f(dx_{11} + br^2\tilde{f}^{-1}d\varphi)^2] , \quad (3.9)$$

$$C_3 = \tilde{b}r^2\tilde{f}^{-1}dx_{11} \wedge dx_9 \wedge d\varphi . \quad (3.10)$$

It reduces to the previous flat background (3.4) when $\tilde{b} = 0$. This $d = 11$ background is related by the 9-11 flip to the background obtained by lifting the NS-NS solution directly to $d = 11$ along x_{11} .

Another special case is $b = 0$, when the metric (3.9) becomes

$$ds_{11}^2 = \tilde{f}^{1/3}(-dt^2 + dx_s^2) + \tilde{f}^{-2/3}(\tilde{f}dr^2 + r^2d\varphi^2 + dx_9^2 + dx_{11}^2) . \quad (3.11)$$

Remarkably, this metric (and $|C_3|$ in (3.10)) are *symmetric* under the “9-11” flip.¹⁷ That means that compactifications along x_{11} or along x_9 lead to the same NS-NS background: the $b = 0$ case of the 2-parameter background (2.2),(2.3), i.e. to (2.8).

By applying T- and S- dualities to (3.7),(3.8) it is easy to construct other similar solutions with mixed magnetic fluxes. Furthermore, it is possible to generalize these $d = 10$ and $d = 11$ solutions to the case of more than two magnetic parameters. Examples of such more general solutions will be presented in section 6.

4. Tachyon instabilities of magnetic backgrounds and type 0 tachyon

As was mentioned in section 2.1, there are two magnetic models (2.16) which are equivalent to type II superstring theory in flat space with fermions obeying *antiperiodic* boundary conditions in the x_9 direction [14]. Modulo periodicity in bR and $\tilde{b}\tilde{R}$ (2.15),(2.26) they are simply ($R = R_9$)

- (a) the model with $\tilde{b} = 0$, $bR = 1$; (b) the model with $\tilde{b}\tilde{R} = 1$, $b = 0$.

They are equivalent (T-dual) as weakly-coupled type II superstring theories compactified on a circle.¹⁸ The direct lift of the corresponding $d = 10$ solutions (2.2),(2.3) to $d = 11$ gives M-theory backgrounds (3.5) and (3.11),(3.10). T-duality equivalence is believed to

¹⁷ This metric is regular: its Ricci scalar is $R = \frac{2\tilde{b}^2}{3(1+\tilde{b}^2r^2)^{7/3}}$, $R_{mn}^2 = 23R^2$, and R_{mnl}^2 has similar behavior.

¹⁸ Note that while the periodicity $bR_9 \rightarrow bR_9 + 2n$ is obvious in the background (2.7), its counterpart $\tilde{b}\tilde{R}_9 \rightarrow \tilde{b}\tilde{R}_9 + 2n$ is not visible at the level of the supergravity background (2.8).

hold also for finite coupling, so one expects that M-theory compactified along x_9, x_{11} in the background (3.5) with $bR = 1$ and in (3.11),(3.10) with $\tilde{b}\tilde{R} = 1$ should be equivalent not only in the weak-coupling limit $R_{11} \rightarrow 0$ but also for finite R_{11} . Thus it is natural to conjecture that M-theory in the background (3.11),(3.10) with $\tilde{b}\tilde{R}_9 = 1$ compactified on the (x_9, x_{11}) torus with $(+, +)$ (periodic) boundary conditions for the fermions is equivalent to M-theory in flat space compactified on 2-torus with the $(-, +)$ fermionic boundary conditions.¹⁹

The M-theory description of $d = 10$ type 0A theory in [8] was based on the $d = 11$ background (3.4) with $bR_{11} = 1$. Now we are able to suggest another description of type 0A string theory – as M-theory in the background (3.11),(3.10) with periodic x_9 and $\tilde{b}\tilde{R}_9 = 1$. The type 0 coupling is $g_{s0} = \frac{R_9}{\sqrt{\alpha'}}$. Because of the 9-11 symmetry of (3.11),(3.10) one may consider also the equivalent model with $\tilde{b}\tilde{R}_{11} = 1$ and $g_{s0} = \frac{R_{11}}{\sqrt{\alpha'}}$.

While the reduction of (3.4) to 10 dimensions gives string theory in R-R F7 background, the reduction of (3.11),(3.10) gives apparently much simpler string theory in NS-NS background ((2.2),(2.3) with $b = 0$) the spectrum of which (in *weakly-coupled* regime) we know. However, the 9-11 flip necessary to relate type II and type 0 theories implies that, e.g., a weakly coupled ($R_{11} \ll R_9$) type II theory is mapped into a strongly coupled ($R_9 \ll R_{11}$) type 0 theory or vice versa. That precludes one from drawing immediate conclusions about the presence or absence of tachyons in type 0 theory in the NS-NS Melvin background directly from the known weakly coupled string spectrum. In particular, the weakly coupled type 0 theory in the NS-NS Melvin background is certainly unstable for any value of magnetic field parameter (because of its own tachyon present already in flat space) but it is expected to become stable at strong coupling and critical magnetic field since it should then become equivalent to a weakly coupled type II string in NS-NS background with a small magnetic field parameter.

According to [8], type IIA and type 0A theories in the $\tilde{b} = 0$ R-R Melvin background (3.6),(3.2) are equivalent, being related by a shift of the R-R field strength parameter: $b_0 = b - R_{11}^{-1}$. The above discussion suggests the following generalization of this relation ($R = R_{11}$):

$$\begin{aligned} \text{Type IIA in } (b, \tilde{b}) \text{ Melvin} &= \text{Type 0A in } (b - R^{-1}, \tilde{b}) \text{ Melvin} \\ &= \text{Type 0A in } (b, \tilde{b} - \tilde{R}^{-1}) \text{ Melvin} . \end{aligned} \tag{4.1}$$

¹⁹ It should be noted that while the background (2.2),(2.3) represents an exact (tree-level) superstring solution, its $d = 11$ lift – the curved $d = 11$ background (3.11),(3.10) may, in principle, be deformed (in contrast to the flat background (3.4)) by higher order M-theory corrections.

It was conjectured in [7] that the type 0A tachyon originates from a massive mode (in twisted sector of Σ -orbifold) in microscopic M-theory, i.e. a mode which is absent in the $d = 11$ supergravity spectrum. Here we will present evidence that in the new M-theory description of type 0A theory – as M-theory in the background (3.11),(3.10) at the critical magnetic field $\tilde{b}\tilde{R}_9 = 1$ – the type 0A tachyon corresponds to a particular fluctuation mode of $d = 11$ supergravity multiplet. Below we shall investigate the appearance of instabilities in these models, looking at the effective field theory spectrum in different regions of the parameter space.

4.1. Spectrum of $d = 11$ fluctuations in magnetic backgrounds

As we have seen in section 2.2, the leading tachyonic instability in the model (2.1) with $b = 0$ is associated with a “non-stringy” state and so can be found directly in the effective low-energy field theory, i.e. in the $d = 10$ supergravity. The same tachyonic state in the T-dual $\tilde{b} = 0$ model is the winding state (2.20) and thus it can not be seen in the supergravity approximation. In the corresponding $d = 11$ description this string state should be represented by a winding membrane state. In the U-dual R-R Melvin model (3.6),(3.2) this tachyon should also be a winding string state (as can be understood using the 9-11 flip).

Following the discussion of section 2.2, let us first study some examples where the presence of tachyonic states can be seen directly at the level of supergravity equations expanded near the magnetic background. Applying U-duality (which is a symmetry of the supergravity equations of motion [32]) to an unstable (say, NS-NS) supergravity background gives, of course, a supergravity solution which is also unstable. This allows one to identify a perturbative tachyon state in the R-R background (3.7),(3.8) with $b = 0, \tilde{b} \neq 0$. Lifting a (potentially) unstable $d = 10$ solution to $d = 11$ gives a (potentially) unstable $d = 11$ supergravity background.

Consider the $d = 11$ background (3.9),(3.10) with $b = 0$, i.e. (3.11). Linearizing the $d = 11$ supergravity equations near this background one can find the corresponding Laplace-type equations for the bosonic fluctuation modes. In general, there should be scalar, “spin 1”, “spin 2” fluctuations (we are referring to angular momentum in the (r, φ) plane).

Consider first the equation for some massless scalar fluctuation ψ , i.e. let us assume, as in (2.29), that there is a fluctuation mode that satisfies

$$\partial_\mu(\sqrt{G}G^{\mu\nu}\partial_\nu)\psi = 0 . \quad (4.2)$$

For the metric (3.11) this becomes

$$\left[-\partial_t^2 + \partial_{x_s}^2 + (1 + \tilde{b}^2 r^2)(\partial_{x_9}^2 + \partial_{x_{11}}^2 + \frac{1}{r^2} \partial_\varphi^2) + \frac{1}{r} \partial_r(r \partial_r) \right] \psi = 0 . \quad (4.3)$$

Setting as in (2.31)

$$\begin{aligned} \psi &= e^{iEt + ip_s x_s + ip_9 x_9 + ip_{11} x_{11} + il\varphi} \eta(r) , \\ p_9 &= \frac{m}{R_9} , \quad p_{11} = \frac{n}{R_{11}} , \quad m, n = 0, \pm 1, \dots , \end{aligned} \quad (4.4)$$

we get

$$\left[-E^2 + p_s^2 + (1 + \tilde{b}^2 r^2)(p_9^2 + p_{11}^2 + \frac{l^2}{r^2}) - \frac{1}{r} \partial_r(r \partial_r) \right] \eta(r) = 0 . \quad (4.5)$$

This equation is of the same form as eq. (2.32), so its spectrum is

$$M^2 \equiv E^2 - p_s^2 = p_9^2 + p_{11}^2 + \tilde{b}^2 l^2 + 2\tilde{b} \sqrt{p_9^2 + p_{11}^2} (l_L + l_R + 1) , \quad (4.6)$$

where $l = l_L - l_R = J$ (cf. (2.35)). In the sector $p_{11} = 0$ the spectrum (4.6) coincides with the ten-dimensional counterpart (2.34) with $b = 0$.

To generalize this spectrum to the case of the spin 1 and spin 2 fluctuations we add (as in section 2.2) the spin contributions S_R, S_L to the orbital angular momentum parts. Indeed, since the $d = 11$ background (3.11) has translational symmetry in x_{11} , the fluctuation modes that do not depend on x_{11} , i.e. have $p_{11} = 0$, must be the same as in case of the corresponding $d = 10$ solution (eqs. (2.2), (2.3) with $b = 0$). But the spectrum of the latter is known, being part of the string theory spectrum in the NS-NS Melvin background (2.10), with its supergravity part given by eq. (2.28) with $b = 0$. Thus the $d = 11$ spectrum should be

$$M^2 = p_9^2 + p_{11}^2 + \tilde{b}^2 J^2 + 2\tilde{b} \sqrt{p_9^2 + p_{11}^2} (l_L + l_R + 1 - S_R + S_L) , \quad (4.7)$$

$$J = l_L - l_R + S_L + S_R .$$

This result should also follow directly from the detailed analysis of the supergravity equations for the graviton – 3-form fluctuation modes. A negative contribution to M^2 appears from the gyromagnetic interaction for the supergravity modes with $l_L = l_R = 0$, and $S_R - S_L = 2$, i.e. with the maximal spin $S_R = 1$ and the minimal spin $S_L = -1$. For such states

$$\begin{aligned} M^2 &= p_9^2 + p_{11}^2 - 2\tilde{b} \sqrt{p_9^2 + p_{11}^2} \\ &= m^2 R_9^{-2} + n^2 R_{11}^{-2} - 2\tilde{b} \sqrt{m^2 R_9^{-2} + n^2 R_{11}^{-2}} . \end{aligned} \quad (4.8)$$

This mass spectrum (4.8) exhibits the presence of tachyons. Without loss of generality, we may assume that $R_{11} < R_9$ (otherwise we would identify x_9 as the eleventh dimension). Increasing $\tilde{b}$ from the zero value (corresponding to the stable flat vacuum), the first tachyon appears for $m = \pm 1$, $l = 0$ and $n = 0$ at $\tilde{b} > \tilde{b}_{\text{cr}}$, with $\tilde{b}_{\text{cr}} = \frac{1}{2R_9}$. This is, of course, the same tachyon that was present in the $d = 10$ model (2.36). The first tachyonic state with non-zero p_{11} will be a state with $n = \pm 1$, $l = 0 = m$, at the critical field $\tilde{b}'_{\text{cr}} = \frac{1}{2R_{11}}$. Since $\tilde{b}_{\text{cr}} < \tilde{b}'_{\text{cr}}$, this instability is physically irrelevant – there should be a vacuum transition at smaller field, i.e. before it could appear.

Similar analysis can be repeated for the supergravity part of the spectrum of M-theory in the background (3.5). However, to establish a correspondence with (4.8) implied by T-duality relation in $d = 10$, here we are to look at the *winding* part of the spectrum, i.e. need to go beyond the supergravity approximation. The result for the relevant sector of the spectrum of the flat $d = 11$ model (3.5) with $b \neq 0$ may be obtained by “uplifting” to eleven dimensions of the spectrum (2.17) of the T-dual NS-NS string model and re-interpreting it as *membrane* spectrum. We will only consider the sector $\hat{N}_R = \hat{N}_L = 0$, $m = 0$, where the spectrum (2.17) is

$$M^2 = \frac{w^2 R_9^2}{\alpha'^2} + b^2 J^2 + \frac{2bwR_9}{\alpha'} (l_L + l_R + 1 - S_R + S_L) . \quad (4.9)$$

Re-written in terms of the M-theory parameters

$$\alpha' = (4\pi^2 R_{11} T_2)^{-1} , \quad T_2 = (2\pi l_P^3)^{-1} , \quad (4.10)$$

where T_2 is the membrane tension and l_P is the eleven dimensional Planck length, eq. (4.9) becomes

$$M^2 = (4\pi^2 w R_9 R_{11} T_2)^2 + b^2 J^2 + 8\pi^2 bw R_9 R_{11} T_2 (l_L + l_R + 1 - S_R + S_L) . \quad (4.11)$$

The first term represents the usual contribution to M^2 coming from a membrane of tension T_2 wrapped on a 2-torus (wound w times around x_9 and once around x_{11}). The spectrum is similar to (4.7), as expected from the T-duality relation between the corresponding ten-dimensional models. The counterpart of the first two terms in (4.7) is now a winding term; the gyromagnetic interaction in (4.7) is traded for an analogous gyromagnetic term where the charge is now the winding number w .

In this way we determine part of the spectrum for the $d = 11$ flat model (3.5) which is relevant for the study of instabilities. The winding membrane with quantum numbers $S_R = 1 = -S_L$, $l_L = l_R = 0$ thus has the mass

$$M^2 = (4\pi^2 w R_9 R_{11} T_2)^2 - 8\pi^2 bw R_9 R_{11} T_2 . \quad (4.12)$$

As the magnetic parameter b is increased from zero, there is a critical value at which M^2 for this state changes sign.

4.2. Instabilities in type 0 and type II theories

As discussed above, the M-theory in the magnetic background (3.9),(3.10) with $bR_{11} = 1, \tilde{b} = 0$ (model (a)) or with

$$b = 0, \quad \tilde{b}\tilde{R}_{11} = \frac{\tilde{b}\alpha'_9}{R_{11}} = \frac{\tilde{b}l_P^3}{2\pi R_9 R_{11}} = 1, \quad (4.13)$$

(model (b)), should describe type 0A string theory with parameters

$$\alpha' = \frac{l_P^3}{2\pi R_{11}}, \quad g_{s0} = \frac{R_{11}}{\sqrt{\alpha'}}. \quad (4.14)$$

In eq. (4.13), the parameter $\alpha'_9 = \frac{l_P^3}{2\pi R_9}$ represents the string scale for type IIA string theory in the magnetic background obtained by dimensional reduction in x_9 of the $d = 11$ solution (3.9) with $b = 0$.

The weakly coupled type 0A theory contains a tachyon. The problem of interest is to identify the corresponding state in the spectrum of M-theory in these backgrounds, and to study its evolution as a function of the magnetic field parameter. It is also of interest to see if other tachyonic instabilities may appear in type 0A theory in a different region of parameter space. We will investigate these issues using the expressions for the spectrum given above in (4.7), (4.11).

The type 0A tachyon state has mass $m_0^2 = -\frac{2}{\alpha'}$, $\alpha' = \frac{l_P^3}{2\pi R_{11}}$. This mode is present at least in the perturbative type 0A spectrum, i.e. for $R_{11} \ll l_P$. The type 0B tachyon has the same mass and is present at least in the region where the type 0B coupling is small, i.e. $R_{11} \ll R_9$.

Now consider the description of type 0A string theory in terms of the model (b) based on (3.11). The symmetry of (4.8) under $R_9 \leftrightarrow R_{11}$ is related to the $SL(2, Z)$ isometry of the background (3.11),(3.10). This background may receive quantum corrections, but it is possible that this symmetry (being a $d = 11$ isometry) may still be present in the quantum-corrected solution. Then the M-theory in this background would (as in the flat space [33]) have the $SL(2, Z)$ symmetry. In the picture of [7], the $SL(2, Z)$ symmetry of type 0B string theory may look unexpected, since the directions x_9 and x_{11} are not equivalent, given different boundary conditions for the fermions. However, it was argued in [7] that the perturbative type II orbifold description of type 0B theory combined with the 9-11 flip requires $SL(2, Z)$ symmetry for consistency.

The present description based on model (b) may be viewed as an additional evidence for the S-duality of type 0B theory, i.e. the symmetry under $g_{s0B} \rightarrow g_{s0B}^{-1}$, with $g_{s0B} =$

R_{11}/R_9 . Indeed, given the background (3.11),(4.13), we are free to make dimensional reduction in x_9 or in x_{11} . For $R_{11} \ll R_9$, one shows (as in section 2) that reduction in x_{11} gives a theory which is equivalent to the free superstring theory with fermions antiperiodic in x_9 . This equivalence is expected to hold for any coupling, in particular, for $R_{11} \gg R_9$. But in this region, by reducing in x_9 , one shows that the theory is, at the same time, equivalent to the free superstring theory with fermions antiperiodic in x_{11} .

Let us explore the $d = 11$ spectrum (4.8) in the region $\tilde{b}\tilde{R} = 1 - \epsilon$, where $\epsilon \rightarrow 0$. Dropping $O(\epsilon)$ term we get

$$M^2 = \frac{m^2}{R_9^2} + \frac{n^2}{R_{11}^2} - \frac{2}{\alpha'} \sqrt{m^2 + \frac{n^2 R_9^2}{R_{11}^2}}. \quad (4.15)$$

The state with $m = 0$ has mass

$$M^2 = \frac{n^2}{R_{11}^2} - \frac{2}{\alpha'} \frac{n R_9}{R_{11}}. \quad (4.16)$$

This state is non-perturbative from the point of view of type 0 theory. There is also a state with $n = 0$, $m = 1$, with

$$M^2 = \frac{1}{R_9^2} - \frac{2}{\alpha'}, \quad (4.17)$$

i.e. with the same mass as the type 0A tachyon in the limit of large R_9 .²⁰ Thus in this description of type 0A string theory as M-theory in the background (3.11),(3.10) it is consistent to identify the type 0A tachyon state with a fluctuation mode of the $d = 11$ supergravity multiplet.

Let us consider now the description of type 0A string theory as M-theory in the background of (3.4) of model (a). Setting $bR_{11} = 1 - \epsilon$, $\epsilon \rightarrow 0$, the spectrum (4.12) becomes

$$M^2 = \frac{w^2 R_9^2}{\alpha'^2} - \frac{2}{\alpha'} \frac{w R_9}{R_{11}}. \quad (4.18)$$

This state may be viewed as ‘‘T-dual’’ of the state (4.16).

In this model (a) description, the type 0A tachyon arises as the T-dual counterpart of (4.17). This can be understood by describing the type 0A theory as in section 2 by setting $bR_9 = 1 - \epsilon$. We then get the state with quantum numbers (2.19) and the mass given by (2.27).

²⁰ This mass formula is applicable for $R_{11} \ll R_9$, $R_{11} \ll l_P$, or for $R_9 \ll R_{11}$, $R_9 \ll l_P$, given that the condition (4.13) is symmetric under the interchange of R_9 and R_{11} .

For a general b , this state has the mass given in (2.20). Using the duality relation (4.1), we can consider the correction to the mass of the tachyon state in type 0A theory when a magnetic field b_0 is turned on

$$M^2 = \frac{R_9^2}{\alpha'^2} - \frac{2}{\alpha'}(1 - b_0 R_9) . \quad (4.19)$$

As expected, this is negative for small values of the magnetic field, but becomes positive for $b_0 R_9 > 1 - \frac{R_9^2}{2\alpha'}$. In particular, type 0A theory with $b_0 R_9 \rightarrow 1$ becomes type IIA string theory, and eq. (4.19) shows that the tachyon is absent.

The above formulas are valid in the perturbative regime, where the radius of the coordinate playing the role of the eleventh dimension is much smaller than the Planck length l_P . The membrane interpretation (4.11) of the spectrum is suitable for extensions beyond perturbation theory. Let us comment on how this can be used to support the conjecture of [7] that in the flat-space type 0 theory, the tachyon becomes massive at strong coupling. In string theory in NS-NS Melvin background with $bR = 1$, the negative term in the mass formula comes from a normal ordering constant (see section 2.1 and (2.27)). Let us assume that the same should be true in the supermembrane theory. In the case of fermions obeying antiperiodic boundary conditions on x_{11} , the normal ordering constant in membrane theory can be computed in the large radius limit, $R_9, R_{11} \gg l_P$ [34]. Then, instead of (4.11), one obtains ($w = 1$):

$$M^2 = (4\pi^2 R_9 R_{11} T_2)^2 - 8T_2 R_{11}^2 R_9^2 \sum_{(n,n') \neq (0,0)} [1 - (-1)^n] (n^2 R_{11}^2 + n'^2 R_9^2)^{-\frac{3}{2}} . \quad (4.20)$$

The infinite sum can be expanded in the region $R_{11} \gg R_9$, where M^2 reduces to (2.27), and in the region $R_{11} \ll R_9$, where one gets

$$M^2 = (4\pi^2 R_9 R_{11} T_2)^2 - 28\zeta(3) T_2 \frac{R_9^2}{R_{11}} \left[1 + O\left(\frac{R_{11}^2}{R_9^2}\right) \right] , \quad (4.21)$$

or, in terms of type 0A parameters (4.14),

$$M^2 \cong \frac{R_9^2}{\alpha'^2} \left(1 - \frac{7\zeta(3)}{\pi^2} \frac{1}{g_{s0}^2} \right) . \quad (4.22)$$

This expression (which applies for $R_9^2 \gg \alpha' g_{s0}^2$) suggests, that instead of the tachyon at weak coupling we indeed get a state with $M^2 > 0$ for $g_{s0}^2 > \frac{7\zeta(3)}{\pi^2}$.

4.3. Comment on M-theory at finite temperature

M-theory at finite temperature T may be studied by considering the $d = 11$ theory in euclidean target space with periodic time x_0 , and periodic coordinate x_{11} , and with fermions antiperiodic in x_0 . This picture was recently used in [34] to compute the one-loop free energy in the supergravity limit, and to determine a critical temperature $T = T(g_s)$ which interpolates between the Hagedorn temperature in the $d = 10$ limit $g_s \ll 1$ and $T \sim l_P^{-1}$ in the $d = 11$ limit $g_s \gg 1$.

Since the boundary conditions are similar, we can thus relate M-theory at finite temperature to M-theory compactifications based on the models (a) and (b), with x_0 playing the role of x_9 and $T = (2\pi R_0)^{-1}$. The $SL(2, Z)$ symmetry of the model (b) compactification, if maintained at the full quantum level, would imply that the exact free energy in M-theory should have the symmetry

$$F(g_{\text{eff}}, A, l_P) = g_{\text{eff}} F(g_{\text{eff}}^{-1}, A, l_P) , \quad (4.23)$$

where

$$g_{\text{eff}} = \frac{R_{11}}{R_0} = 2\pi\sqrt{\alpha'} T g_s , \quad A = R_0 R_{11} . \quad (4.24)$$

In terms of string-theory parameters g_s, T, α' this gives

$$T^{-1} F(g_s, T, \alpha') = \bar{T}^{-1} F(\bar{g}_s, \bar{T}, \bar{\alpha}') , \quad (4.25)$$

where

$$\bar{T} = (2\pi g_s \sqrt{\alpha'})^{-1} , \quad \bar{\alpha}' = \alpha' \frac{R_{11}}{R_0} = \alpha' (2\pi \sqrt{\alpha'} T g_s) , \quad \bar{g}_s = \frac{g_s}{(2\pi T \sqrt{\alpha'} g_s)^{\frac{3}{2}}} . \quad (4.26)$$

Thus the duality relates the low and high temperature regimes, as well as the weak and strong coupling regions.

While the bosonic field contributions to the one-loop $d = 11$ supergravity free energy must have this symmetry [34], it is not obvious why it should be present in the full theory with fermions. Our point here was to note that the symmetry (4.25) of the finite temperature M-theory free energy is related to the conjectured $SL(2, Z)$ symmetry of the type 0B string theory.

5. Type 0A tachyon mass shift in $d = 10$ R-R magnetic flux background

Starting with M-theory in the background (3.4) with antiperiodic fermions in x_{11} , corresponding to type 0A theory in the background of the R-R magnetic flux 7-brane (3.6), one has the opposite situation to the one in the case of type IIA theory in the same R-R background [8]: while $b_0 R_{11} = 0$ represents the flat type 0A theory, $b_0 R_{11} = 1$ should give a theory with periodic fermions in the eleventh dimension, i.e. type IIA theory in flat space. This implies that type 0A string theory in the R-R 7-brane background (3.6),(3.2) must become tachyon-free (i.e. the type 0A tachyon should get positive M^2) for a sufficiently large R-R magnetic field.

By using the perturbative correspondence between type 0A theory and type IIA theory (see sections 1 and 2), and interpolating between them with the NS-NS magnetic background (2.7), we argued in section 4.2 that the tachyon of type 0A theory (considered as a limit of 9-d interpolating theory) should disappear at some magnetic coupling b_0 (cf. (4.19)). If we start with the $d = 10$ type 0A theory and turn on the R-R Melvin background (3.6), (3.2), the tachyon should disappear at some magnetic field $0 < b_0 R_{11} \leq 1$, since at $b_0 R_{11} = 1$ the theory is expected to describe the type IIA theory which does not have a tachyon.

As we have seen in section 4, in the M-theory interpretation of type 0 theory based on (3.4), the tachyon state may be identified with a particular winding membrane (“twisted sector”) state. That means, in particular, that the tachyon mass formula should contain a gyromagnetic interaction term linear in R-R magnetic field, which is responsible for changing the sign of M^2 . From the point of view of perturbative type 0 theory, the tachyon is neutral with respect to the R-R field, so such linear coupling must have a non-perturbative origin.²¹

One may still wonder if it is possible to find some indication of the effect of shifting of the tachyon mass by the R-R flux already at the level of perturbative type 0A theory. In this section we shall consider the equation for the type 0A tachyon in the R-R Melvin background (3.6),(3.2) and show that, as in [3,4], the R-R flux shifts the value of tachyon

²¹ This is in contrast to the 9-d reduction case, where momentum and winding tachyon modes carry charges with respect to the NS-NS vector fields (see eq. (2.23) and below). This is related to the following question: if the type 0 tachyon may indeed be interpreted as a winding mode of M-theory orbifold, then it should be complex, not real as it is in perturbative type 0 theory. A natural conjecture then is that the tachyon should get a scalar partner and thus become charged under the R-R vector field in non-perturbative regime.

(mass)² by a positive term, *quadratic* in the field. This gives a complementary argument in support of the conjecture [8] that the tachyon should disappear at sufficiently strong magnetic R-R field.

Our starting point will be the effective action of type 0A theory for the tachyon T from (NS−,NS−) sector and massless fields: the (NS+,NS+) fields (G_{mn}, B_{mn}, ϕ) , the R-R fields (A_m, A_{mnk}) and their ‘doubles’ $(\bar{A}_m, \bar{A}_{mnk})$ from the twisted sector of type IIB orbifold. All type 0 tree amplitudes which involve only the fields from the (NS+,NS+) and (R+,R+) (or (NS+,NS+) and (R−,R−)) sectors are identical to those in the type II theory – in spite of the absence of space-time supersymmetry, the world-sheet supersymmetry implies the same restrictions on the tree-level string effective action as in the type II theory (e.g., the absence of α' corrections to 3-point functions) [3]. The leading second-derivative terms for the NS-NS fields (G_{mn}, B_{mn}, ϕ) fields and T thus have the standard form

$$S_1 = -2 \int d^D x \sqrt{G} e^{-2\phi} \left[R + 4(\partial\phi)^2 - \frac{1}{12} H_{mnk}^2 - \frac{1}{4} (\partial T)^2 - \frac{1}{4} m_0^2 T^2 \right], \quad (5.1)$$

where $m_0^2 = -\frac{2}{\alpha'}$ is the tachyon mass.²² The leading R-R terms in the action of type 0A theory are [3]

$$S_2 = \int d^D x \sqrt{G} \left[(1 + T^2)(|F_2|^2 + |\bar{F}_2|^2 + |F_4|^2 + |\bar{F}_4|^2) + |F_2 \bar{F}_2| T + |F_4 \bar{F}_4| T \right], \quad (5.2)$$

where $|F_n \bar{F}_n| \equiv \frac{1}{n!} F^{m_1 \dots m_n} \bar{F}_{m_1 \dots m_n}$, $F_n = dA_n$. This action may be diagonalized by introducing the “electric” and “magnetic” combinations $F_{n\pm} = \frac{1}{\sqrt{2}}(F_n \pm \bar{F}_n)$.

It is possible to embed any solution of type II theory into type 0 theory by setting the twisted sector fields to zero: $\bar{F}_n = 0$, $T = 0$ (for $T = 0$ there will be no non-trivial mixing between the R-R forms from the standard and “twisted” sectors) [3,4]. In particular, the R-R 7-brane background (3.6), (3.2) is also a solution ($F_{mn} \neq 0$, $\bar{F}_{mn} = 0$) of type 0A string theory.

The equation for the type 0A tachyon fluctuation in the type 0A counterpart of type IIA R-R vector background follows from the action (5.1),(5.2)

$$S = -2 \int d^{10} x \sqrt{-G} \left(e^{-2\phi} \left[R + 4(\partial\phi)^2 - \frac{1}{4} (\partial T)^2 - \frac{1}{4} m_0^2 T^2 \right] - \frac{1}{4} F_{mn} F^{mn} (1 + \frac{1}{2} T^2) \right). \quad (5.3)$$

²² The world-sheet supersymmetry also constrains the tachyonic sector of the action: all NS-NS amplitudes involving *odd* powers of T (e.g., T^3) vanish. One is able to show also that there is no $T^2 H_{mnk}^2$ term in the action [35]. For a recent discussion of type 0 actions and their T-duality properties see [36].

It follows from here that (part of) the effect that switching on R-R magnetic flux shifts the tachyon (mass)² in positive direction appears to be visible already in the weakly coupled type 0A theory: the tachyon mass is indeed shifted by a term quadratic in the magnetic R-R flux

$$-\frac{1}{\sqrt{G}e^{-2\phi}}\partial_m(\sqrt{G}e^{-2\phi}G^{mn}\partial_n T) + m_{\text{eff}}^2 T = 0, \quad (5.4)$$

$$m_{\text{eff}}^2 = m_0^2 + \frac{1}{2}F_{mn}^2 e^{2\phi}. \quad (5.5)$$

This perturbative equation may not be, of course, sufficient to argue that the spectrum of the scalar T fluctuations in the R-R Melvin background (3.6), (3.2) becomes non-tachyonic for sufficiently large field. As follows from (3.2), the dilaton (effective string coupling) blows up at large r , and thus the above tree-level $d = 10$ effective action is not applicable there.

We may try, however, to concentrate at the near-core (small r) region where the weakly-coupled type 0 string description should be valid.²³ The maximum of $m_{\text{eff}}^2(r)$ in (5.5) computed using (3.6), (3.2) (here b_0 denotes the value of b in type 0 theory)

$$m_{\text{eff}}^2 = m_0^2 + \frac{4b_0^2 e^{2\phi_0}}{(1 + b_0^2 r^2)^{5/2}} \quad (5.6)$$

is reached at $r = 0$

$$m_{\text{eff}}^2(0) = -\frac{2}{\alpha'} + 4b_0^2 g_{s0}^2, \quad g_{s0} = e^{\phi_0}. \quad (5.7)$$

Assuming that the string coupling $g_{s0} = \frac{R_{11}}{\sqrt{\alpha'}}$ is small, we would conclude from (5.7) that the critical value of the magnetic field parameter at which the tachyon becomes effectively massless near $r = 0$ is $\sqrt{2\alpha'}b_*g_{s0} = 1$. It is clear that α' corrections may become important here: since we are assuming $g_{s0} < 1$, we get $\alpha'b_*^2 > 1$.²⁴

Still, it is possible that α' corrections do not qualitatively change the structure of the expression for the mass (5.6), (5.7) so that this effective field theory result is indeed an indication that the tachyon mass is shifted by strong magnetic R-R field.²⁵ As already mentioned at the beginning of this section, non-perturbative corrections are expected to modify (5.7) further, inducing a term linear in the R-R magnetic field.

²³ Since we are going to consider $r \rightarrow 0$, i.e., in particular, $r \sim \sqrt{\alpha'}$, we should be prepared to include all α' corrections.

²⁴ Indeed, the scalar curvature $R(r) = -\frac{3b^2(4+3b_0^2r^2)}{(1+b_0^2r^2)^{5/2}}$, which reaches its maximum at $r = 0$ ($R(0) = -12b^2$), is then strong in string units: $\alpha'R > 1$.

²⁵ A similar effect of shifting of the type 0B tachyon mass by the R-R 5-form flux was observed in [3,4]. There the dilaton was constant, so the theory was weakly coupled everywhere in space, and it was found that one is to require that the scale of the $AdS_5 \times S^5$ solution, i.e. $L = (4\pi g_s N)^{1/4} \sqrt{\alpha'}$, should be of order $\sqrt{\alpha'}$, i.e. sufficiently small, to change the sign of the tachyon mass squared.

6. A class of magnetic flux-brane solutions in $d = 11$

By applying T and S duality transformations to the “seven-brane” solution (3.7), (3.8) one can generate other similar flux brane backgrounds. For example, T-dualities along the x_s directions give “flux p-brane” (or Fp-brane) solutions with $p < 6$. The case of $p = 4$ or F4-brane solution is

$$ds_{10A}^2 = f^{1/2} \left[-dt^2 + dx_1^2 + dx_2^2 + dx_3^2 + dx_4^2 + f^{-1}(dx_5^2 + dx_6^2) + \tilde{f}^{-1}dx_9^2 + dr^2 + r^2 f^{-1} \tilde{f}^{-1} d\varphi^2 \right],$$

$$e^{2(\phi - \phi_0)} = f^{\frac{1}{2}} \tilde{f}^{-1}, \quad A_{\varphi 56} = br^2 f^{-1}, \quad B_{\varphi 9} = \tilde{b} r^2 \tilde{f}^{-1}. \quad (6.1)$$

This is the dimensional reduction along x_{11} of the following $d = 11$ solution

$$ds_{11}^2 = (f\tilde{f})^{1/3} \left[-dt^2 + dx_1^2 + dx_2^2 + dx_3^2 + dx_4^2 + f^{-1}(dx_5^2 + dx_6^2) + \tilde{f}^{-1}(dx_9^2 + dx_{11}^2) + dr^2 + r^2 f^{-1} \tilde{f}^{-1} d\varphi^2 \right], \quad (6.2)$$

$$C_{\varphi 56} = br^2 f^{-1}, \quad C_{\varphi 911} = \tilde{b} r^2 \tilde{f}^{-1}. \quad (6.3)$$

Notice a remarkable symmetry of this background under the interchange of f and $\tilde{f}$ or b and $\tilde{b}$, combined with relabelling of the coordinates $(5, 6) \leftrightarrow (9, 11)$. This is another 11-d counterpart (in addition to (3.9), (3.10)) of our basic 2-parameter magnetic background (2.2), (2.3).

In fact, this is a particular case of a more general solution with four independent magnetic field parameters²⁶

$$ds_{11}^2 = (f_1 f_2 f_3 f_4)^{1/3} \left[-dt^2 + f_1^{-1}(dx_1^2 + dx_2^2) + f_2^{-1}(dx_3^2 + dx_4^2) + f_3^{-1}(dx_5^2 + dx_6^2) + f_4^{-1}(dx_7^2 + dx_8^2) + dr^2 + r^2 f_1^{-1} f_2^{-1} f_3^{-1} f_4^{-1} d\varphi^2 \right], \quad (6.4)$$

$$C_{\varphi 12} = b_1 r^2 f_1^{-1}, \quad C_{\varphi 34} = b_2 r^2 f_2^{-1},$$

$$C_{\varphi 56} = b_3 r^2 f_3^{-1}, \quad C_{\varphi 78} = b_4 r^2 f_4^{-1}, \quad f_i = 1 + b_i^2 r^2. \quad (6.5)$$

²⁶ Note that this metric has a structure similar to that of the background describing supersymmetric [37] intersection of four M5-branes [38]. Dimensional reduction of this background to $d = 4$ gives a 4-parameter generalization of the Melvin solution constructed in [39], where also a more general class of composite $d = 4$ black hole in magnetic field solutions was found (these are somewhat different from $d = 4$ reductions of backgrounds discussed below in section 7).

This solution may be obtained by starting with (6.2),(6.3) and mixing φ with other “parallel” coordinates. To explain this construction in more detail let us first rederive (6.2),(6.3). Let us start with flat $d = 11$ Minkowski space with $C_3 = 0$, and make a formal coordinate shift $\varphi \rightarrow \varphi + b_1 x_8$ as in (3.4). By dimensional reduction in x_8 and T-duality in x_1, x_2 , we get a type IIA solution with the R-R 3-form having the $A_{\varphi 12}$ component. Uplifting it back to eleven dimensions, this gives the background (6.4) with $b_2 = b_3 = b_4 = 0$, i.e. (6.2) with $\tilde{b} = 0$ or (3.11). Next, we start with this non-trivial $d = 11$ background and repeat the same procedure: replace $\varphi \rightarrow \varphi + b_2 x_2$, and make dimensional reduction in x_2 . We obtain a type IIA solution with A_φ and $B_{\varphi 1}$ fields. By T-duality in x_3, x_4 , and uplift, we get (6.2), or (6.4) with $b_3 = b_4 = 0$. Iterating this procedure, we introduce b_3 and b_4 and finally get (6.4),(6.5). This background can be further generalized by making a shift $\varphi \rightarrow \varphi + b'_i x_i$, leading to magnetic solutions in 10 dimensions with more free magnetic parameters.

Finally, let us mention another example of $d = 11$ solution which is obtained by uplifting to 11 dimensions the electric/magnetic F2-brane

$$\begin{aligned}
ds_{11}^2 &= \tilde{f}^{1/3} f^{2/3} \left[-dt^2 + dx_1^2 + dx_2^2 + f^{-1}(dx_3^2 + dx_4^2 + dx_5^2 + dx_6^2) \right. \\
&\quad \left. + \tilde{f}^{-1} dx_9^2 + f^{-1} \tilde{f}^{-1} dx_{11}^2 + dr^2 + r^2 f^{-1} \tilde{f}^{-1} d\varphi^2 \right] , \\
C_{t12} &= br^2 f^{-1} , \quad C_{\varphi 911} = \tilde{b} r^2 \tilde{f}^{-1} .
\end{aligned} \tag{6.6}$$

Interchanging t and, say, x_3 , gives another magnetic solution.

All such $d = 11$ magnetic flux brane solutions are non-supersymmetric and (like the background (3.11),(3.10)) may be perturbatively unstable in certain region of parameter space.

7. Superpositions of magnetic flux branes and D-branes

So far we described some brane-type solutions with magnetic fluxes: instead of magnetic fluxes through spheres (which describe magnetic charges), they have axially symmetric magnetic fluxes through planes. Other classes of flux-brane solutions were presented in [16,40,41].

It is of some interest to investigate “hybrid” solutions which have both brane charges and Melvin-type magnetic fluxes. One way to construct them is to start with M5-brane or M2-brane solution in $d = 11$, pick up a 2-plane (in parallel or transverse directions) with

the metric $dx_1^2 + dx_2^2 = dr^2 + r^2 d\varphi^2$, and then shift $\varphi \rightarrow \varphi + bx_{11}$. Dimensional reduction along x_{11} (combined with U-dualities) then generates a superposition of a D-brane with the K-K Melvin magnetic flux through the (x_1, x_2) plane.

The twist $\varphi \rightarrow \varphi + bx_{11}$ breaks supersymmetry of the original M-brane background. As in the case of the absence of M-brane charges, the magnetic parameter b allows one to smoothly interpolate between periodic and antiperiodic boundary conditions for the fermions. Since all solutions of type II string theory may be embedded into type 0 string theory (see section 5), and in view of the (perturbative and non-perturbative) relations between type II and type 0 theories discussed above, this magnetic b -interpolation may allow one to relate D-branes in type II theory to D-branes in type 0 theory.

For example, consider a type IIA Dp-brane superposed with an axially symmetric magnetic flux with parameter b . While for small string coupling or small bR_{11} the $d = 10$ type IIA description is applicable, this is no longer so for $bR_{11} \sim 1$ (the effective 11-d radius becomes large), i.e. one can no longer use the supergravity approximation based on the dimensionally reduced ($d = 10$) solution. The description of the $bR_{11} \sim 1$ case should be based on M-theory with fermions being antiperiodic along x_{11} . Following [8,16], one may then conjecture that the result should be equivalent to a Dp-brane in type 0 theory in a R-R Melvin background with a small magnetic field with $b_0 R_{11} = 1 - bR_{11}$.

Below we shall present examples of such “hybrid” solutions with a hope that they may be useful in future studies of connections between D-branes of type II and type 0 theories.

Starting with a $d = 11$ M-brane solution and mixing the angular coordinate φ in the (x_1, x_2) plane with x_{11} to add magnetic flux, one has several options: x_1, x_2, x_{11} may be parallel directions to M-brane; x_1, x_2 may be parallel while x_{11} may be transverse; x_{11} may be parallel while x_1, x_2 may be transverse, etc. There are also several options of making dimensional reduction and applying dualities to generate various $d = 10$ “hybrid” brane backgrounds. Let us list some of the solutions that can be obtained in this way. We will use the notation $H_n = 1 + \frac{L^n}{\rho^n}$ for the harmonic function of an M-brane (ρ is the radial direction transverse to the brane).

From the M5-brane background with a twist in the parallel 2-plane in the metric

$$ds_{11}^2 = H_3^{-1/3}[-dt^2 + dx_3^2 + dx_4^2 + dx_{11}^2 + dr^2 + r^2(d\varphi + bdx_{11})^2] + H_3^{2/3}[d\rho^2 + \rho^2 d\Omega_4^2], \quad (7.1)$$

one obtains the D4-brane with a R-R magnetic flux on the world-volume:

$$ds_{10A}^2 = f^{1/2} \left(H_3^{-1/2}[-dt^2 + dx_3^2 + dx_4^2 + dr^2 + r^2 f^{-1} d\varphi^2] + H_3^{1/2}[d\rho^2 + \rho^2 d\Omega_4^2] \right), \quad (7.2)$$

$$e^{2\phi} = H_3^{-1/2}(\rho)f^{3/2}(r) , \quad A_\varphi = br^2f^{-1} , \quad f(r) = 1 + b^2r^2 . \quad (7.3)$$

T-duality then gives a (smeared version of) D3-brane solution with R-R 2-form flux,

$$ds_{10B}^2 = f^{1/2} \left(H_3^{-1/2} [-dt^2 + dx_3^2 + dr^2 + r^2 f^{-1} d\varphi^2] + H_3^{1/2} [f^{-1} dx_4^2 + d\rho^2 + \rho^2 d\Omega_4^2] \right) , \quad (7.4)$$

$$e^{2\phi} = f(r) , \quad A_{\varphi 4} = br^2 f^{-1}(r) .$$

Starting with the M5-brane with the “parallel” twist (7.1) smeared along x_{11}

$$ds_{11}^2 = H_2^{-1/3} [-dt^2 + dx_1^2 + dx_2^2 + dx_3^2 + dr^2 + r^2 (d\varphi + b dx_{11})^2] + H_2^{2/3} [dx_{11}^2 + d\rho^2 + \rho^2 d\Omega_3^2] , \quad (7.5)$$

one obtains the NS5-brane with R-R Melvin magnetic flux:

$$ds_{10A}^2 = \hat{f}^{1/2} \left(-dt^2 + dx_3^2 + dx_4^2 + dx_5^2 + dr^2 + r^2 \hat{f}^{-1} d\varphi^2 + H_2 [d\rho^2 + \rho^2 d\Omega_3^2] \right) , \quad (7.6)$$

$$e^{2\phi} = H_2 \hat{f}^{3/2} , \quad A_\varphi = br^2 H_2^{-1} \hat{f}^{-1} , \quad \hat{f}(r, \rho) \equiv 1 + \frac{b^2 r^2}{H_2(\rho)} , \quad H_2 = 1 + \frac{L^2}{\rho^2} .$$

The smeared M5-brane with two extra isometries with transverse twist

$$ds_{11}^2 = H_1^{-1/3} [-dt^2 + dx_3^2 + \dots + dx_6^2 + dx_{11}^2] + H_1^{2/3} [dr^2 + r^2 (d\varphi + b dx_{11})^2 + d\rho^2 + \rho^2 d\Omega_2^2] , \quad (7.7)$$

gives the D4-brane with R-R magnetic flux in transverse space

$$ds_{10A}^2 = \hat{f}^{1/2} \left(H_1^{-1/2} [-dt^2 + dx_3^2 + \dots + dx_6^2] + H_1^{1/2} [dr^2 + r^2 \hat{f}^{-1} d\varphi^2 + d\rho^2 + \rho^2 d\Omega_2^2] \right) , \quad (7.8)$$

$$e^{2\phi} = H_1^{-1/2} \hat{f}^{3/2} , \quad A_\varphi = br^2 H_1 \hat{f}^{-1} , \quad \hat{f}(r, \rho) \equiv 1 + b^2 r^2 H_1(\rho) .$$

From the M2-brane with three extra isometries, one obtains a D2-brane solution with transverse R-R magnetic flux:

$$ds_{10A}^2 = f^{1/2} \left(H_3^{-1/2} [-dt^2 + dx_3^2 + dx_4^2] + H_3^{1/2} [dr^2 + r^2 f^{-1} d\varphi^2 + d\rho^2 + \rho^2 d\Omega_4^2] \right) , \quad (7.9)$$

$$e^{2\phi} = H_3^{1/2} f^{3/2} , \quad A_\varphi = br^2 f^{-1} .$$

One may apply a similar procedure of a twist and dimensional reduction to the 11-d plane wave background, getting a superposition of D0-brane and R-R flux 7-brane. Here the R-R 1-form will have both electric and magnetic components.

Another way to construct “hybrid” solutions is to start with conformal NS-NS sigma model describing NS5-brane and to replace the 5 free parallel directions by another conformal theory – the 2-parameter magnetic model (2.1). The resulting background has the following metric and dilaton [42]

$$ds_{10A}^2 = -dt^2 + dx_1^2 + dx_2^2 + dr^2 + \tilde{f}^{-1} [r^2 (d\varphi + b dx_9)^2 + dx_9^2] + H_2 (d\rho^2 + \rho^2 d\Omega_3^2) , \quad (7.10)$$

$$e^{2\phi} = e^{2\phi_0} H_2 \tilde{f}^{-1} , \quad \tilde{f} = 1 + \tilde{b}^2 r^2 , \quad H_2 = 1 + \frac{L^2}{\rho^2} , \quad (7.11)$$

as well as NS-NS 2-form which is a combination the one in (2.3) with NS5-brane one, i.e.

$$dB_2 = \frac{2\tilde{b}r}{(1 + \tilde{b}^2 r^2)^2} dr \wedge d\varphi \wedge dx_9 + L^2 d\text{Vol}(S^3) . \quad (7.12)$$

Acknowledgements

We are grateful to S. Frolov for very useful discussions. The work of A.A.T. is partially supported by the DOE grant DE-FG02-91ER40690, PPARC SPG grant 00613, INTAS project 991590 and CRDF Award RPI-2108. Part of this work was done while A.A.T. was participating in the M-theory program at ITP, Santa Barbara supported by the NSF grant PHY99-07949.

Appendix A. Constant magnetic field background

For completeness, let us mention also another example of a magnetic background which in a sense comes closer than Melvin solution to the aim of modelling a uniform magnetic field in closed string theory.

In general relativity, magnetic flux lines gravitationally attract each other and thus tend to concentrate near $r = 0$. A way to have a homogeneous (covariantly constant) magnetic field is to add a rotation: intuitively, it is clear that magnetic flux lines which would otherwise cluster near $r = 0$ would then repel due to centrifugal forces. Such NS-NS background was discussed in [26,43] by giving a magnetic Kaluza-Klein interpretation to the following gravitational wave background (related to a non-semisimple WZW model [44])

$$ds_{10A}^2 = -dt^2 + dx_s^2 + dx_9^2 + 2br^2 d\varphi(dx_9 - dt) + dr^2 + r^2 d\varphi^2 , \quad (A.1)$$

$$B_2 = br^2 d\varphi \wedge (dt - dx_9) , \quad b = \text{const} . \quad (A.2)$$

As was shown in [26,43] the corresponding conformal string model is exactly solvable in terms of free fields and has Hamiltonian similar to (2.10). A more general 3-parameter class of solvable models interpolating between (2.1) and (A.1) was discussed in [18] (see also [20]).

Applying U-duality transformations to (A.1),(A.2) one can construct related supergravity solutions with R-R magnetic fields. One way to do that is first to lift the above background to 11 dimensions and then reduce down along a different spatial direction. The $d = 11$ solution corresponding to (A.1),(A.2) is

$$ds_{11}^2 = -dt^2 + dx_s^2 + dx_9^2 + 2br^2 d\varphi(dx_9 - dt) + dr^2 + r^2 d\varphi^2 + dx_{11}^2 , \quad (\text{A.3})$$

$$C_3 = br^2 dx_{11} \wedge d\varphi \wedge (dt - dx_9) . \quad (\text{A.4})$$

Dimensional reduction in the x_9 direction then gives ($x \equiv x_{11}$)

$$ds_{10A}^2 = -(dt + br^2 d\varphi)^2 + dr^2 + r^2 d\varphi^2 + dx_s^2 + dx^2 , \quad (\text{A.5})$$

$$B_2 = br^2 d\varphi \wedge dx , \quad A_1 = br^2 d\varphi , \quad A_3 = br^2 d\varphi \wedge dt \wedge dx . \quad (\text{A.6})$$

A special property of these backgrounds is that the gauge field strength and curvature invariants are *constant*.

The NS-NS model (A.1) has tachyonic modes with high spin for arbitrarily small values of the magnetic field [43]. It would be interesting to study the stability of the R-R model (A.5),(A.6).

References

- [1] L. J. Dixon and J. A. Harvey, “String Theories In Ten-Dimensions Without Space-Time Supersymmetry,” Nucl. Phys. B **274**, 93 (1986). N. Seiberg and E. Witten, “Spin Structures In String Theory,” Nucl. Phys. B **276**, 272 (1986). C. Thorn, unpublished.
- [2] A. A. Tseytlin, “Sigma model approach to string theory effective actions with tachyons,” J. Math. Phys. **42**, 2854 (2001) [hep-th/0011033].
- [3] I. R. Klebanov and A. A. Tseytlin, “D-branes and dual gauge theories in type 0 strings,” Nucl. Phys. B **546**, 155 (1999) [hep-th/9811035].
- [4] I. R. Klebanov and A. A. Tseytlin, “A non-supersymmetric large N CFT from type 0 string theory,” JHEP **9903**, 015 (1999) [hep-th/9901101].
- [5] I. R. Klebanov, “Tachyon stabilization in the AdS/CFT correspondence,” Phys. Lett. B **466**, 166 (1999) [hep-th/9906220].
- [6] A. Adams and E. Silverstein, “Closed string tachyons, AdS/CFT, and large N QCD,” hep-th/0103220.
- [7] O. Bergman and M. R. Gaberdiel, “Dualities of type 0 strings,” JHEP **9907**, 022 (1999) [hep-th/9906055].
- [8] M. S. Costa and M. Gutperle, “The Kaluza-Klein Melvin solution in M-theory,” hep-th/0012072.
- [9] M.A. Melvin, Phys. Lett. B **8**, 65 (1964).
- [10] G. W. Gibbons and K. Maeda, “Black Holes And Membranes In Higher Dimensional Theories With Dilaton Fields,” Nucl. Phys. B **298**, 741 (1988).
- [11] G.W. Gibbons, “Quantized Flux Tubes In Einstein-Maxwell Theory And Noncompact Internal Spaces,” in: *Fields and Geometry*, Proceedings of the 22nd Karpacz Winter School of Theoretical Physics, ed. A. Jadczyk (World Scientific, Singapore, 1986).
- [12] F. Dowker, J. P. Gauntlett, G. W. Gibbons and G. T. Horowitz, “The Decay of magnetic fields in Kaluza-Klein theory,” Phys. Rev. D **52**, 6929 (1995) [hep-th/9507143]. F. Dowker, J. P. Gauntlett, S. B. Giddings and G. T. Horowitz, “On pair creation of extremal black holes and Kaluza-Klein monopoles,” Phys. Rev. D **50**, 2662 (1994) [hep-th/9312172]. F. Dowker, J. P. Gauntlett, D. A. Kastor and J. Traschen, “Pair creation of dilaton black holes,” Phys. Rev. D **49**, 2909 (1994) [hep-th/9309075].
- [13] F. Dowker, J. P. Gauntlett, G. W. Gibbons and G. T. Horowitz, “Nucleation of P -Branes and Fundamental Strings,” Phys. Rev. D **53**, 7115 (1996) [hep-th/9512154].
- [14] J. G. Russo and A. A. Tseytlin, “Magnetic flux tube models in superstring theory,” Nucl. Phys. B **461**, 131 (1996) [hep-th/9508068].
- [15] J. G. Russo and A. A. Tseytlin, “Green-Schwarz superstring action in a curved magnetic Ramond-Ramond background,” JHEP **9804**, 014 (1998) [hep-th/9804076].
- [16] M. Gutperle and A. Strominger, “Fluxbranes in string theory,” hep-th/0104136.

- [17] A. A. Tseytlin, “Melvin solution in string theory,” *Phys. Lett. B* **346**, 55 (1995) [hep-th/9411198].
- [18] J. G. Russo and A. A. Tseytlin, “Exactly solvable string models of curved space-time backgrounds,” *Nucl. Phys. B* **449**, 91 (1995) [hep-th/9502038].
- [19] A. A. Tseytlin, “Closed superstrings in magnetic field: instabilities and supersymmetry breaking”, *Nucl. Phys. Proc. Suppl.* **49**, 338 (1996) [hep-th/9510041].
- [20] A. A. Tseytlin, “Exact solutions of closed string theory,” *Class. Quant. Grav.* **12**, 2365 (1995) [hep-th/9505052].
- [21] J. D. Blum and K. R. Dienes, “Strong/weak coupling duality relations for non-supersymmetric string theories,” *Nucl. Phys. B* **516**, 83 (1998) [hep-th/9707160].
M. Dine, P. Huet and N. Seiberg, “Large And Small Radius In String Theory,” *Nucl. Phys. B* **322**, 301 (1989).
- [22] R. Rohm, “Spontaneous Supersymmetry Breaking In Supersymmetric String Theories,” *Nucl. Phys. B* **237**, 553 (1984).
- [23] J. J. Atick and E. Witten, “The Hagedorn Transition And The Number Of Degrees Of Freedom Of String Theory,” *Nucl. Phys. B* **310**, 291 (1988).
- [24] Y. Imamura, “Branes in type 0 / type II duality,” *Prog. Theor. Phys.* **102**, 859 (1999) [hep-th/9906090].
B. Craps and F. Roose, “NS fivebranes in type 0 string theory,” *JHEP* **9910**, 007 (1999) [hep-th/9906179].
- [25] M. B. Green, “Wilson-Polyakov loops for critical strings and superstrings at finite temperature,” *Nucl. Phys. B* **381**, 201 (1992).
- [26] J. G. Russo and A. A. Tseytlin, “Constant magnetic field in closed string theory: An Exactly solvable model,” *Nucl. Phys. B* **448**, 293 (1995) [hep-th/9411099].
G. T. Horowitz and A. A. Tseytlin, “A New class of exact solutions in string theory,” *Phys. Rev. D* **51**, 2896 (1995) [hep-th/9409021].
- [27] C. Kounnas and B. Rostand, “Coordinate Dependent Compactifications And Discrete Symmetries,” *Nucl. Phys. B* **341**, 641 (1990).
- [28] A. Abouelsaood, C. G. Callan, C. R. Nappi and S. A. Yost, “Open Strings In Background Gauge Fields,” *Nucl. Phys. B* **280**, 599 (1987).
- [29] N. Seiberg and E. Witten, “String theory and noncommutative geometry,” *JHEP* **9909**, 032 (1999), hep-th/9908142.
- [30] N. K. Nielsen and P. Olesen, “An Unstable Yang-Mills Field Mode,” *Nucl. Phys. B* **144**, 376 (1978).
- [31] E. Bergshoeff, E. Sezgin and P. K. Townsend, “Properties Of The Eleven-Dimensional Super Membrane Theory,” *Annals Phys.* **185**, 330 (1988).
- [32] C. M. Hull and P. K. Townsend, “Unity of superstring dualities,” *Nucl. Phys. B* **438**, 109 (1995) [hep-th/9410167].
- [33] J. H. Schwarz, “An $SL(2, \mathbb{Z})$ multiplet of type IIB superstrings,” *Phys. Lett. B* **360**, 13 (1995) [hep-th/9508143].

- [34] J.G. Russo, “Free energy and critical temperature in eleven dimensions”, Nucl. Phys. B **602**, 109 (2001) [hep-th/0101132].
- [35] A. A. Tseytlin, “Type 0 strings and gauge theories,” *Contributed to 14th International Workshop on High Energy Physics and Quantum Field Theory* (QFTHEP 99), Moscow, June 1999. (http://theory.sinp.msu.ru/~qfthep/99_/Proceed99.html)
- [36] P. Meessen and T. Ortin, “Type 0 T-duality and the tachyon coupling,” hep-th/0103244.
- [37] G. Papadopoulos and P. K. Townsend, “Intersecting M-branes,” Phys. Lett. B **380**, 273 (1996) [hep-th/9603087].
- [38] A. A. Tseytlin, “Harmonic superpositions of M-branes,” Nucl. Phys. B **475**, 149 (1996) [hep-th/9604035].
- [39] R. Emparan, “Composite black holes in external fields,” Nucl. Phys. B **490**, 365 (1997) [hep-th/9610170].
- [40] P. M. Saffin, “Gravitating fluxbranes,” gr-qc/0104014.
- [41] C. Chen, D. V. Gal’tsov and S. A. Sharakin, “Intersecting M-fluxbranes,” Grav. Cosmol. **5**, 45 (1999) [hep-th/9908132].
- [42] G. Papadopoulos, J. G. Russo and A. A. Tseytlin, “Curved branes from string dualities,” Class. Quant. Grav. **17**, 1713 (2000), hep-th/9911253.
- [43] J. G. Russo and A. A. Tseytlin, “Heterotic strings in uniform magnetic field,” Nucl. Phys. B **454**, 164 (1995) [hep-th/9506071].
- [44] C. R. Nappi and E. Witten, “A WZW model based on a nonsemisimple group,” Phys. Rev. Lett. **71**, 3751 (1993) [hep-th/9310112].